

File 2. References of studies included in the database

1. A-Tjak JGL, Morina N, Topper M, Emmelkamp PMG. A Randomized Controlled Trial in Routine Clinical Practice Comparing Acceptance and Commitment Therapy with Cognitive Behavioral Therapy for the Treatment of Major Depressive Disorder. *Psychotherapy and psychosomatics*. 2018;(no pagination).
2. Aagaard J, Foldager L, Makki A, Hansen V, Møller-Nielsen K. The efficacy of psychoeducation on recurrent depression: a randomized trial with a 2-year follow-up. *Nordic journal of psychiatry* [Internet]. 2017; 71(3):[223-9 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/pdf/10.1080/08039488.2016.1266385?needAccess=true>
3. Abas M, Nyamayaro P, Bere T, Saruchera E, Mthobi N, Simms V, et al. Feasibility and acceptability of a task-shifted intervention to enhance adherence to HIV medication and improve depression in people living with HIV in Zimbabwe, a low income country in sub-Saharan Africa. *AIDS and behavior*. 2018;22(1):86-101.
4. Abdollahi A, LeBouthillier D, Najafi M, Asmundson G, Hosseinian S, Shahidi S, et al. Effect of exercise augmentation of cognitive behavioural therapy for the treatment of suicidal ideation and depression. *Journal of affective disorders* [Internet]. 2017; 219:[58-63 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1399-1601.2017.01403.1.x>
5. Agyapong V, Juhas M, Ohinmaa A, Omeje J, Mrklas K, Suen V, et al. Randomized controlled pilot trial of supportive text messages for patients with depression. *BMC psychiatry* [Internet]. 2017; 17(1) (no pagination). Available from: <http://onlinelibrary.wiley.com/doi/pdf/10.1186/s12888-017-1448-2>
6. Ahmadpanah M, Nazaribadie M, Aghaei E, Ghaleiha A, Bakhtiari A, Haghighi M, et al. Influence of adjuvant detached mindfulness and stress management training compared to pharmacologic treatment in primiparae with postpartum depression. *Archives of women's mental health* [Internet]. 2017:[1-9 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/pdf/10.1007/s00737-017-0753-6>
7. Ahmadpanah M, Paghale SJ, Bakhtyari A, Kaikhavani S, Aghaei E, Nazaribadie M, et al. Effects of psychotherapy in combination with pharmacotherapy, when compared to pharmacotherapy only on blood pressure, depression, and anxiety in female patients with hypertension. *Journal of Health Psychology*. 2016;21(7):1216-27.
8. Alexopoulos G, Raue P, McCulloch C, Kanellopoulos D, Seirup J, Sirey J, et al. Clinical Case Management versus Case Management with Problem-Solving Therapy in Low-Income, Disabled Elders with Major Depression: A Randomized Clinical Trial. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry* [Internet]. 2016; 24(1):[50-9 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/pdf/10.1111/gps.12140>
9. Alexopoulos GS, Raue P, Arean P. Problem-solving therapy versus supportive therapy in geriatric major depression with executive dysfunction. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2003;11(1):46-52.
10. Alladin A, Alibhai A. Cognitive hypnotherapy for depression: an empirical investigation. *The International journal of clinical and experimental hypnosis*. 2007;55(2):147-66.
11. Allart-Van Dam E, Hosman CMH, Hoogduin CAL, Schaap CPDR. The coping with depression course: Short-term outcomes and mediating effects of a randomized controlled trial in the treatment of subclinical depression. *Behavior Therapy*. 2003;34(3):381-96.
12. Altamura M, Iuso S, Terrone G, Balzotti A, Carnevale R, Malerba S, et al. Comparing interpersonal counseling and antidepressant treatment in primary care patients with anxious and nonanxious major depression disorder: A randomized control trial. *Clinical Neuropsychiatry*. 2017;14(4):257-62.
13. Ammerman RT, Putnam FW, Altaye M, Stevens J, Teeters AR, Van Ginkel JB. A clinical trial of in-home CBT for depressed mothers in home visitation. *Behavior therapy*. 2013;44(3):359-72.
14. Andersson G, Bergström J, Hollnagel E, Carlbring P, Kaldö V, Ekselius L. Internet-based self-help for depression: randomised controlled trial. *British journal of psychiatry* [Internet]. 2005; 187:[456-61 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1399-1601.2005.01572.x>

15. Andersson G, Hesser H, Veilord A, Svedling L, Andersson F, Sleman O, et al. Randomised controlled non-inferiority trial with 3-year follow-up of internet-delivered versus face-to-face group cognitive behavioural therapy for depression. *Journal of affective disorders*. 2013;151(3):986-94.
16. Anguera JA, Gunning FM, Areán PA. Improving late life depression and cognitive control through the use of therapeutic video game technology: A proof-of-concept randomized trial. *and Anxiety*. 2017;34(6):508-17.
17. Appleby L, Warner R, Whitton A, Faragher B. A controlled study of fluoxetine and cognitive-behavioural counselling in the treatment of postnatal depression. *Bmj*. 1997;314(7085):932-6.
18. Arean PA, Perri MG, Nezu AM, Schein RL, Christopher F, Joseph TX. Comparative effectiveness of social problem-solving therapy and reminiscence therapy as treatments for depression in older adults. *Journal of consulting and clinical psychology*. 1993;61(6):1003-10.
19. Arean PA, Raue P, Mackin RS, Kanellopoulos D, McCulloch C, Alexopoulos GS. Problem-solving therapy and supportive therapy in older adults with major depression and executive dysfunction. *The American journal of psychiatry*. 2010;167(11):1391-8.
20. Arjadi R, Nauta MH, Scholte WF, Hollon SD, Chowdhary N, Suryani AO, et al. Internet-based behavioural activation with lay counsellor support versus online minimal psychoeducation without support for treatment of depression: a randomised controlled trial in Indonesia. *The lancet psychiatry*. 2018;5(9):707-16.
21. Ashman T, Cantor JB, Tsaousides T, Spielman L, Gordon W. Comparison of cognitive behavioral therapy and supportive psychotherapy for the treatment of depression following traumatic brain injury: A randomized controlled trial. *Journal of Head Trauma Rehabilitation*. 2014;29(6):467-78.
22. Ashouri A, Atef-Vahid MK, Gharaee B, Rasoulmian M. Effectiveness of meta-cognitive and cognitive-behavioral therapy in patients with major depressive disorder. *Iranian Journal of Psychiatry and Behavioral Sciences*. 2013;7(2):24-34.
23. Ayen I, Hautzinger M. Cognitive behavior therapy for depression in menopausal women. A controlled, randomized treatment study. *Zeitschrift fur Klinische Psychologie und Psychotherapie*. 2004;33(4):290-9.
24. Ayoughi S, Missmahl I, Weierstall R, Elbert T. Provision of mental health services in resource-poor settings: a randomised trial comparing counselling with routine medical treatment in North Afghanistan (Mazar-e-Sharif). *BMC psychiatry*. 2012;12:14.
25. Azevedo Cardoso T, Mondin TC, Spessato BC, Avila Quevedo L, Mattos Souza LD, Silva RA, et al. The impact of anxious symptoms in the remission of depressive symptoms in a clinical trial for depression: Follow-up of six months. *Journal of affective disorders* [Internet]. 2014; 168:[331-6 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1360-0443.2009.02757.x/abstract>
http://ac.els-cdn.com/S0165032714001451/1-s2.0-S0165032714001451-main.pdf?_tid=a9f472e4-d2f6-11e4-b674-00000aacb361&acdnat=1427291938_827b86909e47c26ed7b01e3ebe78696f
http://ac.els-cdn.com/S0165032714001451/1-s2.0-S0165032714001451-main.pdf?_tid=ef34b388-5865-11e5-9b0f-00000aab0f26&acdnat=1441963233_ea8c985b56d1c9bc3fa99979cb000b9b
26. Azhar MZ, Varma SL. Religious psychotherapy in depressive patients. *Psychotherapy and psychosomatics*. 1995;63(3-4):165-8.
27. Bailey E, Stevens A, Larocca M, Scogin F. A randomized controlled trial of a therapeutic intervention for nursing home residents with dementia and depressive symptoms. *Journal of applied gerontology* [Internet]. 2017; 36(7):[895-908 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1177/0733464815627956>
<http://journals.sagepub.com/doi/pdf/10.1177/0733464815627956>
28. Baker AL, Kavanagh DJ, Kay-Lambkin FJ, Hunt SA, Lewin TJ, Carr VJ, et al. Randomized controlled trial of cognitive-behavioural therapy for coexisting depression and alcohol problems: short-term outcome. *Addiction (Abingdon, England)* [Internet]. 2010; 105(1):[87-99 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1360-0443.2009.02757.x/abstract>
<http://onlinelibrary.wiley.com/doi/10.1111/j.1360-0443.2009.02757.x/asset/j.1360-0443.2009.02757.x.pdf?v=1&t=ieffmmnh&s=7178713dc9ff0aa05faae1c9dded6b559d11159a>
29. Barber JP, Barrett MS, Gallop R, Rynn MA, Rickels K. Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. *The Journal of clinical psychiatry*. 2012;73(1):66-73.
30. Barker WA, Scott J, Eccleston D. The Newcastle chronic depression study: results of a treatment regime. *International clinical psychopharmacology*. 1987;2(3):261-72.
31. Barnhofer T, Crane C, Hargus E, Amarasinghe M, Winder R, Williams JM. Mindfulness-based cognitive therapy as a treatment for chronic depression: A preliminary study. *Behaviour research and therapy*. 2009;47(5):366-73.

32. Barrera M. An evaluation of a brief group therapy for depression. *Journal of Consulting and Clinical Psychology*. 1979;47(2):413.
33. Barrett JE, Williams Jr JW, Oxman TE, Frank E, Katon W, Sullivan M, et al. Treatment of dysthymia and minor depression in primary care: A randomized trial in patients aged 18 to 59 years. *Journal of Family Practice*. 2001;50(5):405-12.
34. Bastos AG, Guimaraes LS, Trentini CM. Neurocognitive changes in depressed patients in psychodynamic psychotherapy, therapy with fluoxetine and combination therapy. *Journal of affective disorders*. 2013;151(3):1066-75.
35. Beach SRH, O'Leary KD. Treating depression in the context of marital discord: Outcome and predictors of response of marital therapy versus cognitive therapy. *Behavior Therapy*. 1992;23(4):507-28.
36. Beck AT, Hollon SD, Young JE, Bedrosian RC, Budenz D. Treatment of depression with cognitive therapy and amitriptyline. *Archives of General Psychiatry*. 1985;42(2):142-8.
37. Bedard M, Felteau M, Marshall S, Cullen N, Gibbons C, Dubois S, et al. Mindfulness-based cognitive therapy reduces symptoms of depression in people with a traumatic brain injury: Results from a randomized controlled trial. *Journal of Head Trauma Rehabilitation*. 2014;29(4):E13-E22.
38. Bedford LA, Dietch JR, Taylor DJ, Boals A, Zayfert C. Computer-Guided Problem-Solving Treatment for Depression, PTSD, and Insomnia Symptoms in Student Veterans: a Pilot Randomized Controlled Trial. *Behavior therapy*. 2018;49(5):756-67.
39. Bedi N, Chilvers C, Churchill R, Dewey M, Duggan C, Fielding K, et al. Assessing effectiveness of treatment of depression in primary care. Partially randomised preference trial. *The British journal of psychiatry : the journal of mental science*. 2000;177:312-8.
40. Beeber LS, Holditch-Davis D, Perreira K, Schwartz TA, Lewis V, Blanchard H, et al. Short-term in-home intervention reduces depressive symptoms in Early Head Start Latina mothers of infants and toddlers. *Research in nursing & health*. 2010;33(1):60-76.
41. Beevers C, Pearson R, Hoffman J, Foulser A, Shumake J, Meyer B. Effectiveness of an internet intervention (Deprexis) for depression in a united states adult sample: a parallel-group pragmatic randomized controlled trial. *Journal of consulting and clinical psychology [Internet]*. 2017; 85(4):[367-80 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/399/CN-01413399/frame.html>.
42. Beevers CG, Clasen PC, Enock PM, Schnyer DM. Attention bias modification for major depressive disorder: Effects on attention bias, resting state connectivity, and symptom change. *Journal of abnormal psychology [Internet]*. 2015; 124(3):[463-75 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/958/CN-01090958/frame.html>.
43. Bellack AS, Hersen M, Himmelhoch J. Social skills training compared with pharmacotherapy and psychotherapy in the treatment of unipolar depression. *American Journal of Psychiatry*. 1981;138(12):1562-7.
44. Bellino S, Zizza M, Rinaldi C, Bogetto F. Combined treatment of major depression in patients with borderline personality disorder: a comparison with pharmacotherapy. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2006;51(7):453-60.
45. Bellino S, Zizza M, Rinaldi C, Bogetto F. Combined therapy of major depression with concomitant borderline personality disorder: comparison of interpersonal and cognitive psychotherapy. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2007;52(11):718-25.
46. Berger T, Himmerli K, Gubser N, Andersson G, Caspar F. Internet-based treatment of depression: a randomized controlled trial comparing guided with unguided self-help. *Cognitive behaviour therapy [Internet]*. 2011; 40(4):[251-66 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/193/CN-00851193/frame.html>
<http://www.tandfonline.com/doi/pdf/10.1080/16506073.2011.616531>.
47. Berger T, Krieger T, Sude K, Meyer B, Maercker A. Evaluating an e-mental health program ("deprexis") as adjunctive treatment tool in psychotherapy for depression: results of a pragmatic randomized controlled trial. *Journal of affective disorders [Internet]*. 2017; 227:[455-62 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/207/CN-01432207/frame.html>.
48. Berger T, Krieger T, Sude K, Meyer B, Maercker A. Evaluating an e-mental health program ("deprexis") as adjunctive treatment tool in psychotherapy for depression: results of a pragmatic randomized controlled trial. *Journal of affective disorders*. 2018;227:455-62.
49. Berking M, Ebert D, Cuijpers P, Hofmann SG. Emotion regulation skills training enhances the efficacy of inpatient cognitive behavioral therapy for major depressive disorder: a randomized controlled trial. *Psychotherapy and psychosomatics*. 2013;82(4):234-45.
50. Beutel ME, Weissflog G, Leuteritz K, Wiltink J, Haselbacher A, Ruckes C, et al. Efficacy of short-term psychodynamic psychotherapy (STPP) with depressed breast cancer patients: Results of a randomized

- controlled multicenter trial. *Annals of oncology* [Internet]. 2014; 25(2):[378-84 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1093/annonc/ndt229>
51. Beutler LE, Engle D, Mohr D, Daldrup RJ, Bergan J, Meredith K, et al. Predictors of differential response to cognitive, experiential, and self-directed psychotherapeutic procedures. *Journal of consulting and clinical psychology*. 1991;59(2):333-40.
 52. Beutler LE, Moleiro C, Malik M, Mark Harwood T, Romanelli R, Gallagher-Thompson D, et al. A comparison of the Dodo, EST, and ATI factors among comorbid stimulant-dependent, depressed patients. *Clinical Psychology & Psychotherapy*. 2003;10(2):69-85.
 53. Beutler LE, Scogin F, Kirkish P, Schretlen D, Corbishley A, Hamblin D, et al. Group cognitive therapy and alprazolam in the treatment of depression in older adults. *Journal of consulting and clinical psychology*. 1987;55(4):550-6.
 54. Birney A, Gunn R, Russell J.K., Ary DV. MoodHacker Mobile Web App With Email for Adults to Self-Manage Mild-to-Moderate Depression: Randomized Controlled Trial. *Journal of medical Internet research*. 2016;4(1):2291-5222
 55. Blackburn IM, Bishop S, Glen AIM. The efficacy of cognitive therapy in depression: A treatment trial using cognitive therapy and pharmacotherapy, each alone and in combination. *British Journal of Psychiatry*. 1981;139(3):181-9.
 56. Blackburn IM, Moore RG. Controlled acute and follow-up trial of cognitive therapy and pharmacotherapy in out-patients with recurrent depression. *The British journal of psychiatry : the journal of mental science*. 1997;171:328-34.
 57. Blackwell SE, Browning M, Mathews A, Pictet A, Welch J, Davies J, et al. Positive imagery-based cognitive bias modification as a web-based treatment tool for depressed adults: A randomized controlled trial. *Clinical Psychological Science*. 2015;3(1):91-111.
 58. Blanco C, Markowitz JC, Hellerstein DJ, Nezu AM, Wall M, Olfson M, et al. A randomized trial of interpersonal psychotherapy, problem solving therapy, and supportive therapy for major depressive disorder in women with breast cancer. *Breast cancer research and treatment*. 2018;(no pagination).
 59. Bloch M, Meiboom H, Lorberblatt M, Bluvstein I, Aharonov I, Schreiber S. The effect of sertraline add-on to brief dynamic psychotherapy for the treatment of postpartum depression: a randomized, double-blind, placebo-controlled study. *The Journal of clinical psychiatry*. 2012;73(2):235-41.
 60. Blom K, Jernelöv S, Kraepelien M, Bergdahl MO, Jungmarker K, Ankartjärn L, et al. Internet treatment addressing either insomnia or depression, for patients with both diagnoses: A randomized trial. *Sleep*. 2015;38(2):267-77.
 61. Blom MB, Jonker K, Dusseldorp E, Spinhoven P, Hoencamp E, Haffmans J, et al. Combination treatment for acute depression is superior only when psychotherapy is added to medication. *Psychotherapy and psychosomatics*. 2007;76(5):289-97.
 62. Bodenmann G, Plancherel B, Beach SR, Widmer K, Gabriel B, Meuwly N, et al. Effects of coping-oriented couples therapy on depression: a randomized clinical trial. *Journal of consulting and clinical psychology*. 2008;76(6):944-54.
 63. Boele F, Klein M, Verdonck-de LI, Cuijpers P, Heimans J, Snijders T, et al. Internet-based guided self-help for glioma patients with depressive symptoms: a randomized controlled trial. *Journal of neuro-oncology* [Internet]. 2017:[1-13 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1007/s11060-017-2712-5>
 64. Boele FW, Klein M, Verdonck-de Leeuw IM, Cuijpers P, Heimans JJ, Snijders TJ, et al. Internet-based guided self-help for glioma patients with depressive symptoms: a randomized controlled trial. *Journal of neuro-oncology*. 2018;137(1):191-203.
 65. Boeschoten RE, Dekker J, Uitdehaag BMJ, Beekman ATF, Hoogendoorn AW, Collette EH, et al. Internet-based treatment for depression in multiple sclerosis: A randomized controlled trial. *Multiple Sclerosis*. 2017;23(8):1112-22.
 66. Bohlmeijer ET, Fledderus M, Rokx TA, Pieterse ME. Efficacy of an early intervention based on acceptance and commitment therapy for adults with depressive symptomatology: Evaluation in a randomized controlled trial. *Behaviour research and therapy*. 2011;49(1):62-7.
 67. Bolton P, Bass J, Neugebauer R, Verdelli H, Clougherty KF, Wickramaratne P, et al. Group interpersonal psychotherapy for depression in rural Uganda: a randomized controlled trial. *Jama*. 2003;289(23):3117-24.
 68. Bowers W, Stuart S, Macfarlane R, Gorman L. Use of computer-administered cognitive-behavior therapy with depressed inpatients. *Depression*. 1993;1(6):294-9.
 69. Bowers WA. Treatment of depressed in-patients. Cognitive therapy plus medication, relaxation plus medication, and medication alone. *The British Journal of Psychiatry*. 1990;156(1):73-8.

70. Bowman D, Scogin F, Lyrene B. The efficacy of self-examination therapy and cognitive bibliotherapy in the treatment of mild to moderate depression. *Psychotherapy Research*. 1995;5(2):131-40.
71. Bowman V, Ward LC, Bowman D, Scogin F. Self-examination therapy as an adjunct treatment for depressive symptoms in substance abusing patients. *Addictive behaviors*. 1996;21(1):129-33.
72. Brand E, Clingempeel WG. Group behavioral therapy with depressed geriatric inpatients: An assessment of incremental efficacy. *Behavior Therapy*. 1992;23(3):475-82.
73. Bright JI, Baker KD, Neimeyer RA. Professional and paraprofessional group treatments for depression: a comparison of cognitive-behavioral and mutual support interventions. *Journal of consulting and clinical psychology*. 1999;67(4):491-501.
74. Brown RA, Lewinsohn PM. A psychoeducational approach to the treatment of depression: comparison of group, individual, and minimal contact procedures. *Journal of consulting and clinical psychology*. 1984;52(5):774.
75. Brown RA, Ramsey SE, Kahler CW, Palm KM, Monti PM, Abrams D, et al. A randomized controlled trial of cognitive-behavioral treatment for depression versus relaxation training for alcohol-dependent individuals with elevated depressive symptoms. *Journal of studies on alcohol and drugs*. 2011;72(2):286-96.
76. Brown SA, Glasner-Edwards SV, Tate SR, McQuaid JR, Chalekian J, Granholm E. Integrated cognitive behavioral therapy versus twelve-step facilitation therapy for substance-dependent adults with depressive disorders. *Journal of psychoactive drugs*. 2006;38(4):449-60.
77. Browne G, Steiner M, Roberts J, Gafni A, Byrne C, Dunn E, et al. Sertraline and/or interpersonal psychotherapy for patients with dysthymic disorder in primary care: 6-month comparison with longitudinal 2-year follow-up of effectiveness and costs. *Journal of affective disorders*. 2002;68(2-3):317-30.
78. Buhrman M, Syk M, Burvall O, Hartig T, Gordh T, Andersson G. Individualized guided internet-delivered cognitive-behavior therapy for chronic pain patients with comorbid depression and anxiety: A randomized controlled trial. *Clinical Journal of Pain*. 2015;31(6):504-16.
79. Buntrock C, Ebert D, Lehr D, Riper H, Smit F, Cuijpers P, et al. Effectiveness of a Web-Based Cognitive Behavioural Intervention for Subthreshold Depression: Pragmatic Randomised Controlled Trial. *Psychotherapy and Psychosomatics*. 2015;84(6):348-58.
80. Burnand Y, Andreoli A, Kolatte E, Venturini A, Rosset N. Psychodynamic psychotherapy and clomipramine in the treatment of major depression. *Psychiatric services (Washington, DC)*. 2002;53(5):585-90.
81. Burns A, Banerjee S, Morris J, Woodward Y, Baldwin R, Proctor R, et al. Treatment and prevention of depression after surgery for hip fracture in older people: randomized, controlled trials. *Journal of the American Geriatrics Society*. 2007;55(1):75-80.
82. Burns A, O'Mahen H, Baxter H, Bennert K, Wiles N, Ramchandani P, et al. A pilot randomised controlled trial of cognitive behavioural therapy for antenatal depression. *BMC psychiatry*. 2013;13:33.
83. Butler LD, Waelde LC, Hastings TA, Chen XH, Symons B, Marshall J, et al. Meditation with yoga, group therapy with hypnosis, and psychoeducation for long-term depressed mood: a randomized pilot trial. *Journal of clinical psychology*. 2008;64(7):806-20.
84. Caldwell YT, Steffen PR. Adding HRV biofeedback to psychotherapy increases heart rate variability and improves the treatment of major depressive disorder. *International journal of psychophysiology*. 2018;131:96-101.
85. Carlbring P, Hagglund M, Luthstrom A, Dahlin M, Kadowaki A, Vernmark K, et al. Internet-based behavioral activation and acceptance-based treatment for depression: a randomized controlled trial. *Journal of affective disorders*. 2013;148(2-3):331-7.
86. Carpenter KM, Smith JL, Aharonovich E, Nunes EV. Developing therapies for depression in drug dependence: results of a stage 1 therapy study. *The American journal of drug and alcohol abuse*. 2008;34(5):642-52.
87. Carr A, Finnegan L, Griffin E, Cotter P, Hyland A. A Randomized Controlled Trial of the Say Yes to Life (SYTL) Positive Psychology Group Psychotherapy Program for Depression: An Interim Report. *Journal of Contemporary Psychotherapy*. 2017;47(3):153-61.
88. Carta MG, Petretto D, Adamo S, Bhat KM, Lecca ME, Mura G, et al. Counseling in primary care improves depression and quality of life. *Clinical Practice and Epidemiology in Mental Health*. 2012;8.
89. Carter JD, McIntosh VV, Jordan J, Porter RJ, Frampton CM, Joyce PR. Psychotherapy for depression: a randomized clinical trial comparing schema therapy and cognitive behavior therapy. *Journal of affective disorders*. 2013;151(2):500-5.
90. Casanas R, Catalan R, del Val JL, Real J, Valero S, Casas M. Effectiveness of a psycho-educational group program for major depression in primary care: a randomized controlled trial. *BMC psychiatry*. 2012;12:230.

91. Castillo-Pérez S, Gómez-Pérez V, Calvillo Velasco M, Pérez-Campos E, Mayoral M-A. Effects of music therapy on depression compared with psychotherapy. *The Arts in Psychotherapy*. 2010;37(5):387-90.
92. Castonguay LG, Schut AJ, Aikens DE, Constantino MJ, Laurenceau J-P, Bologh L, et al. Integrative Cognitive Therapy for Depression: A Preliminary Investigation. *Journal of Psychotherapy Integration*. 2004;14(1):4-20.
93. Chan AS, Wong QY, Sze SL, Kwong PP, Han YM, Cheung MC. A Chinese Chan-based mind-body intervention for patients with depression. *Journal of affective disorders*. 2012;142(1-3):283-9.
94. Chan MF, Ng SE, Tien A, Man Ho RC, Thayala J. A randomised controlled study to explore the effect of life story review on depression in older Chinese in Singapore. *Health & social care in the community*. 2013;21(5):545-53.
95. Chaput Y, Magnan A, Gendron A. The co-administration of quetiapine or placebo to cognitive-behavior therapy in treatment refractory depression: a preliminary trial. *BMC psychiatry*. 2008;8:73.
96. Cheavens JS, Strunk DR, Lazarus SA, Goldstein LA. The compensation and capitalization models: a test of two approaches to individualizing the treatment of depression. *Behaviour research and therapy*. 2012;50(11):699-706.
97. Chen CH, Tseng YF, Chou FH, Wang SY. Effects of support group intervention in postnatally distressed women. A controlled study in Taiwan. *Journal of psychosomatic research*. 2000;49(6):395-9.
98. Chesney MA, Chambers DB, Taylor JM, Johnson LM, Folkman S. Coping effectiveness training for men living with HIV: results from a randomized clinical trial testing a group-based intervention. *Psychosomatic medicine*. 2003;65(6):1038-46.
99. Chetty D, Hoque ME. Effectiveness of a nurse facilitated cognitive group intervention among mild to moderately-depressed-women in KwaZulu-Natal, South Africa. *African journal of psychiatry*. 2013;16(1):29-34.
100. Chiang KJ, Chen TH, Hsieh HT, Tsai JC, Ou KL, Chou KR. One-Year Follow-Up of the Effectiveness of Cognitive Behavioral Group Therapy for Patients' Depression: A Randomized, Single-Blinded, Controlled Study. *Scientific World Journal*. 2015;2015.
101. Chibanda D, Shetty AK, Tshimanga M, Woelk G, Stranix-Chibanda L, Rusakaniko S. Group problem-solving therapy for postnatal depression among HIV-positive and HIV-negative mothers in Zimbabwe. *Journal of the International Association of Providers of AIDS Care*. 2014;13(4):335-41.
102. Cho HJ, Kwon JH, Lee JJ. Antenatal cognitive-behavioral therapy for prevention of postpartum depression: a pilot study. *Yonsei medical journal*. 2008;49(4):553-62.
103. Choi I, Zou J, Titov N, Dear BF, Li S, Johnston L, et al. Culturally attuned Internet treatment for depression amongst Chinese Australians: a randomised controlled trial. *Journal of affective disorders [Internet]*. 2012; 136(3):[459-68 pp.]. Available from:
<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2796.2012.01661.x>
http://ac.els-cdn.com/S0165032711007166/1-s2.0-S0165032711007166-main.pdf?_tid=4f864efa-248f-11e5-b688-00000aabb0f27&acdnat=1436263543_a0d3101843d6b73487215995d73d5db2
http://ac.els-cdn.com/S0165032711007166/1-s2.0-S0165032711007166-main.pdf?_tid=fa8db356-586a-11e5-8afc-00000aacb362&acdnat=1441965399_d54359d61f7306c4fa7ee4c72b2819ca
104. Choi NG, Marti CN, Bruce ML, Hegel MT. Depression in homebound older adults: problem-solving therapy and personal and social resourcefulness. *Behavior therapy*. 2013;44(3):489-500.
105. Chowdhary N, Anand A, Dimidjian S, Shinde S, Weobong B, Balaji M, et al. The healthy activity program lay counsellor delivered treatment for severe depression in India: Systematic development and randomised evaluation. *British Journal of Psychiatry*. 2016;208(4):381-8.
106. Choy J, Lou V. Effectiveness of the Modified Instrumental Reminiscence Intervention on Psychological Well-Being Among Community-Dwelling Chinese Older Adults: A Randomized Controlled Trial. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry [Internet]*. 2016; 24(1):[60-9 pp.]. Available from:
<http://onlinelibrary.wiley.com/doi/10.1111/gps.12266>
http://ac.els-cdn.com/S1064748115001670/1-s2.0-S1064748115001670-main.pdf?_tid=95f86b2c-d99f-11e6-9ed6-00000aacb361&acdnat=1484319192_096102ffec244a3a9300739e04d8937f
107. Christensen H, Griffiths KM, Jorm AF. Delivering interventions for depression by using the internet: randomised controlled trial. *Bmj*. 2004;328(7434):265.
108. Christensen H, Griffiths KM, Mackinnon AJ, Brittliffe K. Online randomized controlled trial of brief and full cognitive behaviour therapy for depression. *Psychological medicine*. 2006;36(12):1737-46.
109. Chronis-Tuscano A, Clarke TL, O'Brien KA, Raggi VL, Diaz Y, Mintz AD, et al. Development and preliminary evaluation of an integrated treatment targeting parenting and depressive symptoms in mothers of children with attention-deficit/hyperactivity disorder. *Journal of consulting and clinical psychology*. 2013;81(5):918.

110. Clark R, Pluczek A, Brown R. A mother–infant therapy group model for postpartum depression. *Infant Mental Health Journal*. 2008;29(5):514-36.
111. Clark R, Pluczek A, Wenzel A. Psychotherapy for Postpartum Depression: A Preliminary Report. *American Journal of Orthopsychiatry*. 2003;73(4):441-54.
112. Clarke G, Eubanks D, Reid E, Kelleher C, O'Connor E, DeBar LL, et al. Overcoming Depression on the Internet (ODIN) (2): a randomized trial of a self-help depression skills program with reminders. *Journal of medical Internet research*. 2005;7(2):e16.
113. Clarke G, Kelleher C, Hornbrook M, DeBar L, Dickerson J, Gullion C. Randomized effectiveness trial of an Internet, pure self-help, cognitive behavioral intervention for depressive symptoms in young adults. *Cognitive behaviour therapy*. 2009;38(4):222-34.
114. Clarke G, Reid E, Eubanks D, O'Connor E, DeBar LL, Kelleher C, et al. Overcoming depression on the Internet (ODIN): a randomized controlled trial of an Internet depression skills intervention program. *Journal of medical Internet research*. 2002;4(3):E14.
115. Cohen S, O'Leary KD, Foran H. A randomized clinical trial of a brief, problem-focused couple therapy for depression. *Behavior therapy [Internet]*. 2010; 41(4):[433-46 pp.]. Available from:
<http://onlinelibrary.wiley.com/doi/10.1111/j.1761-9071.2010.01311.x>
http://ac.els-cdn.com/S0005789410000523/1-s2.0-S0005789410000523-main.pdf?_tid=1ab9b3d0-2569-11e5-aeaf-00000aacb35f&acdnat=1436357086_d20d24e7a01aebb9c22ae38e2a516101
http://ac.els-cdn.com/S0005789410000523/1-s2.0-S0005789410000523-main.pdf?_tid=7531412c-586b-11e5-812a-00000aab0f26&acdnat=1441965605_090b80ee8e2f0895a44ba8d35bcacc34
116. Collado A, Calderón M, MacPherson L, Lejuez C. The efficacy of behavioral activation treatment among depressed Spanish-speaking Latinos. *Journal of Consulting and Clinical Psychology*. 2016;84(7):651-7.
117. Constantino MJ, Marnell ME, Haile AJ, Kanther-Sista SN, Wolman K, Zappert L, et al. Integrative cognitive therapy for depression: a randomized pilot comparison. *Psychotherapy (Chicago, Ill) [Internet]*. 2008; 45(2):[122-34 pp.]. Available from:
<http://onlinelibrary.wiley.com/doi/10.1111/j.1761-9071.2008.01311.x>
118. Cooper CL, Hind D, Parry GD, Isaac CL, Dimairo M, O'Cathain A, et al. Computerised cognitive behavioural therapy for the treatment of depression in people with multiple sclerosis: external pilot trial. *Trials*. 2011;12:259.
119. Cooper PJ, Murray L, Wilson A, Romaniuk H. Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I. Impact on maternal mood. *The British journal of psychiatry : the journal of mental science*. 2003;182:412-9.
120. Corruble E, Swartz HA, Bottai T, Vaiva G, Bayle F, Llorca PM, et al. Telephone-administered psychotherapy in combination with antidepressant medication for the acute treatment of major depressive disorder. *Journal of affective disorders*. 2016;190:6-11.
121. Covi L, Lipman RS. Cognitive behavioral group psychotherapy combined with imipramine in major depression. *Psychopharmacology bulletin*. 1987;23(1):173-6.
122. Cramer H, Salisbury C, Conrad J, Eldred J, Araya R. Group cognitive behavioural therapy for women with depression: pilot and feasibility study for a randomised controlled trial using mixed methods. *BMC psychiatry*. 2011;11:82.
123. Daughters SB, Braun AR, Sargeant MN, Reynolds EK, Hopko DR, Blanco C, et al. Effectiveness of a brief behavioral treatment for inner-city illicit drug users with elevated depressive symptoms: the life enhancement treatment for substance use (LETS Act!). *The Journal of clinical psychiatry*. 2008;69(1):122-9.
124. David D, Szentagotai A, Lupu V, Cosman D. Rational emotive behavior therapy, cognitive therapy, and medication in the treatment of major depressive disorder: a randomized clinical trial, posttreatment outcomes, and six-month follow-up. *Journal of clinical psychology*. 2008;64(6):728-46.
125. De Fruyt F, Van Leeuwen K, Bagby RM, Rolland JP, Rouillon F. Assessing and interpreting personality change and continuity in patients treated for major depression. *Psychological assessment*. 2006;18(1):71-80.
126. de Graaf LE, Gerhards SA, Arntz A, Riper H, Metsmakers JF, Evers SM, et al. Clinical effectiveness of online computerised cognitive-behavioural therapy without support for depression in primary care: randomised trial. *The British journal of psychiatry : the journal of mental science*. 2009;195(1):73-80.
127. De Jong M, Peeters F, Gard T, Ashih H, Doorley J, Walker R, et al. A randomized controlled pilot study on mindfulness-based cognitive therapy for unipolar depression in patients with chronic pain. *Journal of clinical psychiatry*. 2018;79(1):26-34.
128. de Jong R, Treiber R, Henrich G. Effectiveness of two psychological treatments for inpatients with severe and chronic depressions. *Cognitive Therapy and Research*. 1986;10(6):645-63.

129. de Jong-Meyer R, Hautzinger M. Results of two multicenter treatment studies among patients with endogenous and nonendogenous depression: Conclusions and prospects. *Zeitschrift fuer Linische Psychologie*. 1996;25(2):155-60.
130. de Jonghe F, Hendricksen M, van Aalst G, Kool S, Peen V, Van R, et al. Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. *The British journal of psychiatry : the journal of mental science*. 2004;185:37-45.
131. de Jonghe F, Kool S, van Aalst G, Dekker J, Peen J. Combining psychotherapy and antidepressants in the treatment of depression. *Journal of affective disorders*. 2001;64(2-3):217-29.
132. de Mello MF, Myczcowisk LM, Menezes PR. A randomized controlled trial comparing moclobemide and moclobemide plus interpersonal psychotherapy in the treatment of dysthymic disorder. *The Journal of psychotherapy practice and research*. 2001;10(2):117-23.
133. Deady M, Mills KL, Teesson M, Kay-Lambkin F. An Online Intervention for Co-Occurring Depression and Problematic Alcohol Use in Young People: Primary Outcomes From a Randomized Controlled Trial. *Journal of medical Internet research*. 2016;18(3):e71.
134. Dekker JJ, Koelen JA, Van HL, Schoevers RA, Peen J, Hendriksen M, et al. Speed of action: the relative efficacy of short psychodynamic supportive psychotherapy and pharmacotherapy in the first 8 weeks of a treatment algorithm for depression. *Journal of affective disorders*. 2008;109(1-2):183-8.
135. Dekker RL, Moser DK, Peden AR, Lennie TA. Cognitive therapy improves three-month outcomes in hospitalized patients with heart failure. *Journal of cardiac failure*. 2012;18(1):10-20.
136. Delgadillo J, Gore S, Ali S, Ekers D, Gilbody S, Gilchrist G, et al. Feasibility randomized controlled trial of cognitive and behavioral interventions for depression symptoms in patients accessing drug and alcohol treatment. *Journal of substance abuse treatment*. 2015;55:6-14.
137. Denton WH, Wittenborn AK, Golden RN. Augmenting antidepressant medication treatment of depressed women with emotionally focused therapy for couples: a randomized pilot study. *Journal of marital and family therapy*. 2012;38 Suppl 1:23-38.
138. DeRubeis RJ, Hollon SD, Amsterdam JD, Shelton RC, Young PR, Salomon RM, et al. Cognitive therapy vs medications in the treatment of moderate to severe depression. *Archives of general psychiatry*. 2005;62(4):409-16.
139. Desautels C, Savard J, Ivers H, Savard MH, Caplette-Gingras A. Treatment of Depressive Symptoms in Patients with Breast Cancer: A Randomized Controlled Trial Comparing Cognitive Therapy and Bright Light Therapy. *Health Psychology*. 2017.
140. Desautels C, Savard J, Ivers H, Savard MH, Caplette-Gingras A. Treatment of depressive symptoms in patients with breast cancer: a randomized controlled trial comparing cognitive therapy and bright light therapy. *Health psychology*. 2018;37(1):1-13.
141. Dimidjian S, Goodman S, Sherwood N, Simon G, Ludman E, Gallop R, et al. A Pragmatic Randomized Clinical Trial of Behavioral Activation for Depressed Pregnant Women. *Journal of consulting and clinical psychology* [Internet]. 2017; 85(1):[26-36 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1037/0893-3200.85.1.26>
142. Dimidjian S, Hollon SD, Dobson KS, Schmaling KB, Kohlenberg RJ, Addis ME, et al. Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression. *Journal of consulting and clinical psychology*. 2006;74(4):658-70.
143. Dindo L, Recober A, Marchman JN, Turvey C, O'Hara MW. One-day behavioral treatment for patients with comorbid depression and migraine: A pilot study. *Behaviour Research and Therapy*. 2012;50(9):537-43.
144. Dobkin RD, Menza M, Allen LA, Gara MA, Mark MH, Tiu J, et al. Cognitive-behavioral therapy for depression in Parkinson's disease: a randomized, controlled trial. *The American journal of psychiatry*. 2011;168(10):1066-74.
145. Doering LV, Chen B, Cross Bodan R, Magsarili MC, Nyamathi A, Irwin MR. Early cognitive behavioral therapy for depression after cardiac surgery. *The Journal of cardiovascular nursing*. 2013;28(4):370-9.
146. Donker T, Bennett K, Bennett A, Mackinnon A, Straten A, Cuijpers P. Internet-delivered interpersonal psychotherapy versus internet-delivered cognitive behavioral therapy for adults with depressive symptoms: Randomized controlled noninferiority trial. *Journal of medical Internet research* [Internet]. 2013; 15(5):[146-61 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1037/1098-9113.15.5.146>
147. Dowrick C, Dunn G, Ayuso-Mateos JL, Dalgard OS, Page H, Lehtinen V. Problem solving treatment and group psychoeducation for depression: multicentre randomised controlled trial Outcomes of Depression International Network [ODIN] Group. *Bmj* [Internet]. 2000; 321(7274):[1450-4 pp.]. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC27549/pdf/1450.pdf>

148. Dozois DJ, Bieling PJ, Patelis-Siotis I, Hoar L, Chudzik S, McCabe K, et al. Changes in self-schema structure in cognitive therapy for major depressive disorder: a randomized clinical trial. *Journal of consulting and clinical psychology*. 2009;77(6):1078-88.
149. Drapkin ML, Tate SR, McQuaid JR, Brown SA. Does initial treatment focus influence outcomes for depressed substance abusers? *Journal of substance abuse treatment*. 2008;35(3):343-50.
150. Driessen E, Van HL, Don FJ, Peen J, Kool S, Westra D, et al. The efficacy of cognitive-behavioral therapy and psychodynamic therapy in the outpatient treatment of major depression: a randomized clinical trial. *The American journal of psychiatry*. 2013;170(9):1041-50.
151. Duarte PS, Miyazaki MC, Blay SL, Sesso R. Cognitive-behavioral group therapy is an effective treatment for major depression in hemodialysis patients. *Kidney international*. 2009;76(4):414-21.
152. Dunlop BW, Kelley ME, Aponte-Rivera V, Mletzko-Crowe T, Kinkead B, Ritchie JC, et al. Effects of patient preferences on outcomes in the Predictors of Remission in Depression to Individual and Combined Treatments (PRedICT) study. *The American journal of psychiatry*. 2017;174(6):546-56.
153. Dunlop BW, Kelley ME, Mletzko TC, Velasquez CM, Craighead WE, Mayberg HS. Depression beliefs, treatment preference, and outcomes in a randomized trial for major depressive disorder. *Journal of psychiatric research*. 2012;46(3):375-81.
154. Dunn NJ, Rehm LP, Schillaci J, Soucek J, Mehta P, Ashton CM, et al. A randomized trial of self-management and psychoeducational group therapies for comorbid chronic posttraumatic stress disorder and depressive disorder. *Journal of traumatic stress*. 2007;20(3):221-37.
155. Dunn RJ. Cognitive modification with depression-prone psychiatric patients. *Cognitive Therapy and Research*. 1979;3(3):307-17.
156. Dunner DL, Schmalting KB, Hendrickson H, Becker J, Lehman A, Bea C. Cognitive therapy versus fluoxetine in the treatment of dysthymic disorder. *Depression*. 1996;4(1):34-41.
157. Dwight-Johnson M, Aisenberg E, Golinelli D, Hong S, O'Brien M, Ludman E. Telephone-based cognitive-behavioral therapy for Latino patients living in rural areas: a randomized pilot study. *Psychiatric services (Washington, DC)*. 2011;62(8):936-42.
158. Ebert DD, Buntrock C, Lehr D, Smit F, Riper H, Baumeister H, et al. Effectiveness of Web- and Mobile-Based Treatment of Subthreshold Depression With Adherence-Focused Guidance: A Single-Blind Randomized Controlled Trial. *Behav Ther*. 2017.
159. Ebert DD, Buntrock C, Lehr D, Smit F, Riper H, Baumeister H, et al. Effectiveness of Web- and Mobile-Based Treatment of Subthreshold Depression With Adherence-Focused Guidance: a Single-Blind Randomized Controlled Trial. *Behavior therapy*. 2018;49(1):71-83.
160. Ebert DD, Lehr D, Baumeister H, Boss L, Riper H, Cuijpers P, et al. GET.ON Mood Enhancer: efficacy of Internet-based guided self-help compared to psychoeducation for depression: an investigator-blinded randomised controlled trial. *Trials*. 2014;15:39.
161. Eddington KM, Silvia PJ, Foxworth TE, Hoet A, Kwapil TR. Motivational deficits differentially predict improvement in a randomized trial of self-system therapy for depression. *Journal of Consulting and Clinical Psychology*. 2015;83(3):602-16.
162. Egede LE, Walker RJ, Payne EH, Knapp RG, Acierno R, Frueh BC. Effect of psychotherapy for depression via home telehealth on glycemic control in adults with type 2 diabetes: subgroup analysis of a randomized clinical trial. *Journal of telemedicine and telecare*. 2018;24(9):596-602.
163. Eisendrath SJ, Gillung E, Delucchi KL, Segal ZV, Nelson JC, McInnes LA, et al. A Randomized Controlled Trial of Mindfulness-Based Cognitive Therapy for Treatment-Resistant Depression. *Psychotherapy and psychosomatics*. 2016;85(2):99-110.
164. Ekeblad A, Falkenström F, Andersson G, Vestberg R, Holmqvist R. Randomized trial of interpersonal psychotherapy and cognitive behavioral therapy for major depressive disorder in a community-based psychiatric outpatient clinic. *Depression and anxiety*. 2016;33(12):1090-8.
165. Ekers D, Richards D, McMillan D, Bland JM, Gilbody S. Behavioural activation delivered by the non-specialist: phase II randomised controlled trial. *The British journal of psychiatry : the journal of mental science*. 2011;198(1):66-72.
166. Ekkers W, Korrelboom K, Huijbrechts I, Smits N, Cuijpers P, Gaag M. Competitive Memory Training for treating depression and rumination in depressed older adults: a randomized controlled trial. *Behaviour research and therapy [Internet]*. 2011; 49(10):[588-96 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1469-7610.2011.02411.x>
http://ac.els-cdn.com/S0005796711001112/1-s2.0-S0005796711001112-main.pdf?_tid=1aad2664-586e-11e5-8445-00000aab0f02&acdnat=1441966742_09aba6f99a1a6f9814cb146c08fee76a.
167. Elkin I, Shea MT, Watkins JT, Imber SD, Sotsky SM, Collins JF, et al. National Institute of Mental Health Treatment of Depression Collaborative Research Program. General effectiveness of treatments. *Archives of general psychiatry*. 1989;46(11):971-82; discussion 83.

168. Emanuels-Zuurveen L, Emmelkamp PM. Individual behavioural-cognitive therapy v. marital therapy for depression in maritally distressed couples. *The British journal of psychiatry : the journal of mental science*. 1996;169(2):181-8.
169. Emanuels-Zuurveen L, Emmelkamp PM. Spouse-aided therapy with depressed patients. *Behavior modification*. 1997;21(1):62-77.
170. Embling S. The effectiveness of cognitive behavioural therapy in depression. *Nursing standard (Royal College of Nursing (Great Britain))* : 1987). 2002;17(14-15):33-41.
171. Eseadi C, Obidoa MA, Ogbuabor SE, Ikechukwu-Ilomuanya AB. Effects of Group-Focused Cognitive-Behavioral Coaching Program on Depressive Symptoms in a Sample of Inmates in a Nigerian Prison. *International journal of offender therapy and comparative criminology*. 2018;62(6):1589-602.
172. Euteneuer F, Dannehl K, Del Rey A, Engler H, Schedlowski M, Rief W. Immunological effects of behavioral activation with exercise in major depression: An exploratory randomized controlled trial. *Translational Psychiatry*. 2017;7(5).
173. Evans RL, Connis RT. Comparison of brief group therapies for depressed cancer patients receiving radiation treatment. *Public health reports (Washington, DC : 1974)*. 1995;110(3):306-11.
174. Fann JR, Bombardier CH, Vannoy S, Dyer J, Ludman E, Dikmen S, et al. Telephone and in-person cognitive behavioral therapy for major depression after traumatic brain injury: a randomized controlled trial. *J Neurotrauma*. 2015;32(1):45-57.
175. Faramarzi M, Alipor A, Esmaelzadeh S, Kheirkhah F, Poladi K, Pash H. Treatment of depression and anxiety in infertile women: cognitive behavioral therapy versus fluoxetine. *Journal of affective disorders*. 2008;108(1-2):159-64.
176. Farrer L, Christensen H, Griffiths KM, Mackinnon A. Internet-based CBT for depression with and without telephone tracking in a national helpline: randomised controlled trial. *PLoS One*. 2011;6(11):e28099.
177. Faulconbridge LF, Driscoll CFB, Hopkins CM, Bailer Benforado B, Bishop-Gilyard C, Carvajal R, et al. Combined Treatment for Obesity and Depression: a Pilot Study. *Obesity (Silver Spring, Md)*. 2018;26(7):1144-52.
178. Feixas G, Bados A, Garcia-Grau E, Paz C, Montesano A, Compan V, et al. A dilemma-focused intervention for depression: a multicenter, randomized controlled trial with a 3-month follow-up. *Depression and anxiety [Internet]*. 2016; 33(9):[862-9 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/da.23200>
179. Field T, Diego M, Delgado J, Medina L. Peer support and interpersonal psychotherapy groups experienced decreased prenatal depression, anxiety and cortisol. *Early human development*. 2013;89(9):621-4.
180. Field T, Diego MA, Hernandez-Reif M, Schanberg S, Kuhn C. Massage therapy effects on depressed pregnant women. *Journal of psychosomatic obstetrics and gynaecology*. 2004;25(2):115-22.
181. Finkenzeller W, Zobel I, Rietz S, Schramm E, Berger M. [Interpersonal psychotherapy and pharmacotherapy for post-stroke depression. Feasibility and effectiveness]. *Der Nervenarzt*. 2009;80(7):805-12.
182. Fischer A, Schroder J, Vettorazzi E, Wolf OT, Pottgen J, Lau S, et al. An online programme to reduce depression in patients with multiple sclerosis: A randomised controlled trial. *The Lancet Psychiatry [Internet]*. 2015; 2(3):[217-23 pp.]. Available from: [http://onlinelibrary.wiley.com/doi/10.1016/S2053-2554\(15\)00010-0](http://onlinelibrary.wiley.com/doi/10.1016/S2053-2554(15)00010-0)
183. Fissler M, Winnebeck E, Schroeter TA, Gumbertsbach M, Huntenburg JM, Gärtner M, et al. Brief training in mindfulness may normalize a blunted error-related negativity in chronically depressed patients. *Cognitive, Affective & Behavioral Neuroscience*. 2017;17(6):1164-75.
184. Fledderus M, Bohlmeijer ET, Pieterse ME, Schreurs KM. Acceptance and commitment therapy as guided self-help for psychological distress and positive mental health: a randomized controlled trial. *Psychological medicine*. 2012;42(3):485-95.
185. Floyd M, Scogin F, McKendree-Smith NL, Floyd DL, Rokke PD. Cognitive therapy for depression: a comparison of individual psychotherapy and bibliotherapy for depressed older adults. *Behavior modification*. 2004;28(2):297-318.
186. Folke F, Hursti T, Tungstrom S, Soderberg P, Kanter JW, Kuutmann K, et al. Behavioral activation in acute inpatient psychiatry: A multiple baseline evaluation. *Journal of behavior therapy and experimental psychiatry [Internet]*. 2015; 46:[170-81 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/jbpe.21200>

187. Folke F, Parling T, Melin L. Acceptance and commitment therapy for depression: A preliminary randomized clinical trial for unemployed on long-term sick leave. *Cognitive and Behavioral Practice*. 2012;19(4):583-94.
188. Fonagy P, Rost F, Carlyle JA, McPherson S, Thomas R, Pasco Fearon RM, et al. Pragmatic randomized controlled trial of long-term psychoanalytic psychotherapy for treatment-resistant depression: The Tavistock Adult Depression Study (TADS). *World Psychiatry*. 2015;14(3):312-21.
189. Forand N, Barnett J, Strunk D, Hindiyeh M, Feinberg J, Keefe J. Efficacy of Guided iCBT for Depression and Mediation of Change by Cognitive Skill Acquisition. *Behavior therapy [Internet]*. 2017; (no pagination). Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/486/CN-01429486/frame.html>.
190. Forand NR, Barnett JG, Strunk DR, Hindiyeh MU, Feinberg JE, Keefe JR. Efficacy of Guided iCBT for Depression and Mediation of Change by Cognitive Skill Acquisition. *Behavior Therapy*. 2018;49(2):295-307.
191. Forsell E, Bendix M, Holländare F, Szymanska von Schultz B, Nasiell J, Blomdahl-Wetterholm M, et al. Internet delivered cognitive behavior therapy for antenatal depression: A randomised controlled trial. *Journal of Affective Disorders*. 2017;221:56-64.
192. Frank E, Cassano GB, Rucci P, Thompson WK, Kraemer HC, Fagiolini A, et al. Predictors and moderators of time to remission of major depression with interpersonal psychotherapy and SSRI pharmacotherapy. *Psychological medicine*. 2011;41(1):151-62.
193. Freedland KE, Carney RM, Rich MW, Steinmeyer BC, Rubin EH. Cognitive behavior therapy for depression and self-care in heart failure patients a randomized clinical trial. *JAMA Internal Medicine*. 2015;175(11):1773-82.
194. Freedland KE, Skala JA, Carney RM, Rubin EH, Lustman PJ, D'vila-Rom-n VG, et al. Treatment of depression after coronary artery bypass surgery: a randomized controlled trial. *Archives of general psychiatry [Internet]*. 2009; 66(4):[387-96 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/578/CN-00687578/frame.html> http://archpsyc.jamanetwork.com/data/Journals/PSYCH/11878/yoa80111_387_396.pdf.
195. Freeman MP, Davis M, Sinha P, Wisner KL, Hibbeln JR, Gelenberg AJ. Omega-3 fatty acids and supportive psychotherapy for perinatal depression: a randomized placebo-controlled study. *Journal of affective disorders*. 2008;110(1-2):142-8.
196. Freire E, Williams C, Messow C-M, Cooper M, Elliott R, McConnachie A, et al. Counselling versus low-intensity cognitive behavioural therapy for persistent sub-threshold and mild depression (CLICD): A pilot/feasibility randomised controlled trial. *BMC psychiatry*. 2015;15.
197. Fremont J, Wilcoxon Craighead L. Aerobic exercise and cognitive therapy in the treatment of dysphoric moods. *Cognitive Therapy and Research*. 1987;11(2):241-51.
198. Friedman AS. Interaction of drug therapy with marital therapy in depressive patients. *Archives of general psychiatry*. 1975;32(5):619-37.
199. Fry P. Structured and unstructured reminiscence training and depression among the elderly. *Clinical Gerontologist*. 1983;1(3):15-37.
200. Furukawa TA, Horikoshi M, Kawakami N, Kadota M, Sasaki M, Sekiya Y, et al. Telephone cognitive-behavioral therapy for subthreshold depression and presenteeism in workplace: a randomized controlled trial. *PLoS One*. 2012;7(4):e35330.
201. Gallagher-Thompson D, Steffen AM. Comparative effects of cognitive-behavioral and brief psychodynamic psychotherapies for depressed family caregivers. *Journal of consulting and clinical psychology*. 1994;62(3):543-9.
202. García-Peña C, Vázquez-Estupiñan F, Avalos-Pérez F, Jiménez LVR, Sánchez-García S, Juárez-Cedillo T. Clinical effectiveness of group cognitive-behavioural therapy for depressed older people in primary care: A randomised controlled trial. *Salud Mental*. 2015;38(1):33-9.
203. Garnefski N, Kraaij V, Schroevers M. Effects of a cognitive behavioral self-help program on depressed mood for people with acquired chronic physical impairments: a pilot randomized controlled trial. *Patient education and counseling*. 2011;85(2):304-7.
204. Gater R, Waheed W, Husain N, Tomenson B, Aseem S, Creed F. Social intervention for British Pakistani women with depression: randomised controlled trial. *The British journal of psychiatry : the journal of mental science*. 2010;197(3):227-33.
205. Gaudio BA, Busch AM, Wenzel SJ, Nowlan K, Epstein-Lubow G, Miller IW. Acceptance-based behavior therapy for depression with psychosis: Results from a pilot feasibility randomized controlled trial. *Journal of Psychiatric Practice*. 2015;21(5):320-33.
206. Gawrysiak M, Nicholas C, Hopko DR. Behavioral activation for moderately depressed university students: Randomized controlled trial. *Journal of counseling psychology [Internet]*. 2009; 56(3):[468-75

- pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/589/CN-00738589/frame.html>.
207. Geisner IM, Varvil-Weld L, Mittmann AJ, Mallett K, Turrisi R. Brief web-based intervention for college students with comorbid risky alcohol use and depressed mood: does it work and for whom? *Addictive behaviors*. 2015;42:36-43.
 208. Gellis ZD, Bruce ML. Problem solving therapy for subthreshold depression in home healthcare patients with cardiovascular disease. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2010;18(6):464-74.
 209. Gellis ZD, McGinty J, Tierney L, Jordan C, Burton J, Misener E. Randomized controlled trial of problem-solving therapy for minor depression in home care. [References]. *Research on social work practice [Internet]*. 2008; 18(6):[596-606 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/563/CN-00738563/frame.html>.
 210. Geraedts AS, Kleiboer AM, Wiezer NM, van Mechelen W, Cuijpers P. Short-term effects of a web-based guided self-help intervention for employees with depressive symptoms: randomized controlled trial. *Journal of medical Internet research*. 2014;16(5):e121.
 211. Gibbons MB, Thompson SM, Scott K, Schauble LA, Mooney T, Thompson D, et al. Supportive-expressive dynamic psychotherapy in the community mental health system: a pilot effectiveness trial for the treatment of depression. *Psychotherapy (Chicago, Ill)*. 2012;49(3):303-16.
 212. Gilbody S, Brabyn S, Lovell K, Kessler D, Devlin T, Smith L, et al. Telephone-supported computerised cognitive-behavioural therapy: REEACT-2 large-scale pragmatic randomised controlled trial. *British journal of psychiatry [Internet]*. 2017; 210(5):[362-7 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/387/CN-01372387/frame.html> <http://bjp.rcpsych.org/content/bjprcpsych/210/5/362.full.pdf>.
 213. Gilbody S, Littlewood E, Hewitt C, Brierley G, Tharmanathan P, Araya R, et al. Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REEACT trial): large scale pragmatic randomised controlled trial. *BMJ*. 2015;351:h5627.
 214. Gitlin LN, Harris LF, McCoy MC, Chernet N, Pizzi LT, Jutkowitz E, et al. A home-based intervention to reduce depressive symptoms and improve quality of life in older African Americans: a randomized trial. *Annals of internal medicine*. 2013;159(4):243-52.
 215. Glozier N, Christensen H, Naismith S, Cockayne N, Donkin L, Neal B, et al. Internet-delivered cognitive behavioural therapy for adults with mild to moderate depression and high cardiovascular disease risks: a randomised attention-controlled trial. *PLoS One*. 2013;8(3):e59139.
 216. Glueckauf RL, Davis WS, Willis F, Sharma D, Gustafson DJ, Hayes J, et al. Telephone-based, cognitive-behavioral therapy for african american dementia caregivers with depression: Initial findings. *Rehabilitation Psychology*. 2012;57(2):124-39.
 217. Gois C, Dias VV, Carmo I, Duarte R, Ferro A, Santos AL, et al. Treatment response in type 2 diabetes patients with major depression. *Clinical psychology & psychotherapy*. 2014;21(1):39-48.
 218. Goldman RN, Greenberg LS, Angus L. The effects of adding emotion-focused interventions to the client-centered relationship conditions in the treatment of depression. *Psychotherapy Research*. 2006;16(5):537-49.
 219. Goodman JH, Prager J, Goldstein R, Freeman M. Perinatal Dyadic Psychotherapy for postpartum depression: a randomized controlled pilot trial. *Archives of women's mental health*. 2015;18(3):493-506.
 220. Greenberg J, Datta T, Shapero BG, Sevinc G, Mischoulon D, Lazar SW. Compassionate hearts protect against wandering minds: Self-compassion moderates the effect of mind-wandering on depression. *Spirituality in Clinical Practice*. 2018;5(3):155-69.
 221. Greenberg LS, Watson J. Experiential therapy of depression: Differential effects of client-centered relationship conditions and process experiential interventions. *Psychotherapy Research*. 1998;8(2):210-24.
 222. Griffiths KM, Mackinnon AJ, Crisp DA, Christensen H, Bennett K, Farrer L. The effectiveness of an online support group for members of the community with depression: a randomised controlled trial. *PLoS one*. 2012;7(12):e53244.
 223. Grote NK, Swartz HA, Geibel SL, Zuckoff A, Houck PR, Frank E. A randomized controlled trial of culturally relevant, brief interpersonal psychotherapy for perinatal depression. *Psychiatric services (Washington, DC)*. 2009;60(3):313-21.
 224. Gumley A, White R, Briggs A, Ford I, Barry S, Stewart C, et al. A parallel group randomised open blinded evaluation of Acceptance and Commitment Therapy for depression after psychosis: pilot trial outcomes (ADAPT). *Schizophrenia research [Internet]*. 2017; 183:[143-50 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/430/CN-01373430/frame.html>.
 225. Hagen R, Hjemdal O, Solem S, Kennair LEO, Nordahl HM, Fisher P, et al. Metacognitive therapy for depression in adults: A waiting list randomized controlled trial with six months follow-up. *Frontiers in Psychology*. 2017;8.

226. Haller M, Norman SB, Cummins K, Trim RS, Xu X, Cui R, et al. Integrated Cognitive Behavioral Therapy Versus Cognitive Processing Therapy for Adults With Depression, Substance Use Disorder, and Trauma. *Journal of substance abuse treatment*. 2016;62:38-48.
227. Hallford D, Mellor D. Autobiographical memory-based intervention for depressive symptoms in young adults: A randomized controlled trial of cognitive-remembrance therapy. *Psychotherapy and psychosomatics* [Internet]. 2016; 85(4):[246-9 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/psy.12177>
<http://www.karger.com/Article/Pdf/444417>.
228. Hallgren M, Kraepelien M, Öjehagen A, Lindefors N, Zeebari Z, Kaldo V, et al. Physical exercise and internet-based cognitive-behavioural therapy in the treatment of depression: Randomised controlled trial. *British Journal of Psychiatry*. 2015;207(3):227-34.
229. Hamamci Z. Integrating psychodrama and cognitive behavioral therapy to treat moderate depression. *Arts in Psychotherapy*. 2006;33(3):199-207.
230. Hamdan-Mansour AM, Puskar K, Bandak AG. Effectiveness of cognitive-behavioral therapy on depressive symptomatology, stress and coping strategies among Jordanian university students. *Issues in mental health nursing*. 2009;30(3):188-96.
231. Haringsma R, Engels G, Cuijpers P, Spinhoven P. Effectiveness of the Coping With Depression (CWD) course for older adults provided by the community-based mental health care system in the Netherlands: a randomized controlled field trial. *International Psychogeriatrics*. 2006;18(02):307-25.
232. Harley R, Sprich S, Safren S, Jacobo M, Fava M. Adaptation of dialectical behavior therapy skills training group for treatment-resistant depression. *The Journal of nervous and mental disease*. 2008;196(2):136-43.
233. Hassiotis A, Serfaty M, Azam K, Strydom A, Blizard R, Romeo R, et al. Manualised Individual Cognitive Behavioural Therapy for mood disorders in people with mild to moderate intellectual disability: A feasibility randomised controlled trial. *Journal of affective disorders*. 2013;151(1):186-95.
234. Hautzinger M, De Jong-Meyer R, Treiber R, Rudolf GAE, Thien U. Efficacy of cognitive behavior therapy, pharmacotherapy, and the combination of both in non-melancholic, unipolar depression. *Zeitschrift für Klinische Psychologie*. 1996;25(2):130-45.
235. Hautzinger M, Welz S. Kognitive Verhaltenstherapie bei Depressionen im Alter: Ergebnisse einer kontrollierten Vergleichsstudie unter ambulanten Bedingungen an Depressionen mittleren Schweregrads. = Cognitive behavioral therapy for depressed older outpatients--A controlled, randomized trial. *Zeitschrift für Gerontologie und Geriatrie*. 2004;37(6):427-35.
236. Hautzinger M, Welz S. Kurz- und längerfristige wirksamkeit psychologischer interventionen bei depressionen im alter. = Short- and long-term efficacy of psychological intervention for depression in older adults. *Zeitschrift für Klinische Psychologie und Psychotherapie: Forschung und Praxis*. 2008;37(1):52-60.
237. Hayden T, Perantie DC, Nix BD, Barnes LD, Mostello DJ, Holcomb WL, et al. Treating prepartum depression to improve infant developmental outcomes: a study of diabetes in pregnancy. *Journal of clinical psychology in medical settings*. 2012;19(3):285-92.
238. Hayman PM, Cope CS. Effects of assertion training on depression. *Journal of clinical psychology*. 1980;36(2):534-43.
239. Heckman TG, Heckman BD, Anderson T, Lovejoy TI, Markowitz JC, Shen Y, et al. Tele-interpersonal psychotherapy acutely reduces depressive symptoms in depressed HIV-infected rural persons: A randomized clinical trial. *Behavioral Medicine*. 2017;43(4):285-95.
240. Heckman TG, Markowitz JC, Heckman BD, Woldu H, Anderson T, Lovejoy TI, et al. A randomized clinical trial showing persisting reductions in depressive symptoms in HIV-infected rural adults following brief telephone-administered interpersonal psychotherapy. *Annals of Behavioral Medicine*. 2018;52(4):299-308.
241. Heckman TG, Sikkema KJ, Hansen N, Kochman A, Heh V, Neufeld S, et al. A randomized clinical trial of a coping improvement group intervention for HIV-infected older adults. *Journal of behavioral medicine*. 2011;34(2):102-11.
242. Hegerl U, Hautzinger M, Mergl R, Kohlen R, Schütze M, Scheunemann W, et al. Effects of pharmacotherapy and psychotherapy in depressed primary-care patients: a randomized, controlled trial including a patients' choice arm. *International journal of neuropsychopharmacology* [Internet]. 2010; 13(1):[31-44 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1532-5415.2009.02881.x>
<http://ijnp.oxfordjournals.org/content/ijnp/13/1/31.full.pdf>.
243. Hellerstein DJ, Little SA, Samstag LW, Batchelder S, Muran JC, Fedak M, et al. Adding group psychotherapy to medication treatment in dysthymia: a randomized prospective pilot study. *The Journal of psychotherapy practice and research*. 2001;10(2):93-103.

244. Hermanns N, Schmitt A, Gahr A, Herder C, Nowotny B, Roden M, et al. The effect of a Diabetes-Specific Cognitive Behavioral Treatment Program (DIAMOS) for patients with diabetes and subclinical depression: results of a randomized controlled trial. *Diabetes care*. 2015;38(4):551-60.
245. Hernandez-Avila CA, Modesto-Lowe V, Feinn R, Kranzler HR. Nefazodone treatment of comorbid alcohol dependence and major depression. *Alcoholism, clinical and experimental research*. 2004;28(3):433-40.
246. Herrmann-Lingen C, Beutel ME, Bosbach A, Deter H-C, Fritzsche K, Hellmich M, et al. A stepwise psychotherapy intervention for reducing risk in coronary artery disease (SPIRR-CAD): Results of an observer-blinded, multicenter, randomized trial in depressed patients with coronary artery disease. *Psychosomatic Medicine*. 2016;78(6):704-15.
247. Himelhoch S, Medoff D, Maxfield J, Dihmes S, Dixon L, Robinson C, et al. Telephone based cognitive behavioral therapy targeting major depression among urban dwelling, low income people living with HIV/AIDS: results of a randomized controlled trial. *AIDS and behavior*. 2013;17(8):2756-64.
248. Hitchcock C, Gormley S, Rees C, Rodrigues E, Gillard J, Panesar I, et al. A randomised controlled trial of memory flexibility training (MemFlex) to enhance memory flexibility and reduce depressive symptomatology in individuals with major depressive disorder. *Behaviour research and therapy*. 2018;110:22-30.
249. Hobfoll SE, Blais RK, Stevens NR, Walt L, Gengler R. Vets prevail online intervention reduces PTSD and depression in veterans with mild-to-moderate symptoms. *Journal of Consulting and Clinical Psychology*. 2016;84(1):31-42.
250. Hoifodt RS, Lillevoll KR, Griffiths KM, Wilsgaard T, Eisemann M, Waterloo K, et al. The clinical effectiveness of web-based cognitive behavioral therapy with face-to-face therapist support for depressed primary care patients: randomized controlled trial. *Journal of medical Internet research*. 2013;15(8):e153.
251. Holden JM, Sagovsky R, Cox JL. Counselling in a general practice setting: Controlled study of health visitor intervention in treatment of postnatal depression. *British Medical Journal*. 1989;298(6668):223-6.
252. Hollon SD, DeRubeis RJ, Evans MD, Wiemer MJ, Garvey MJ, Grove WM, et al. Cognitive therapy and pharmacotherapy for depression. Singly and in combination. *Archives of general psychiatry*. 1992;49(10):774-81.
253. Hollon SD, DeRubeis RJ, Fawcett J, Amsterdam JD, Shelton RC, Zajecka J, et al. Effect of cognitive therapy with antidepressant medications vs antidepressants alone on the rate of recovery in major depressive disorder: a randomized clinical trial. *JAMA psychiatry*. 2014;71(10):1157-64.
254. Honey KL, Bennett P, Morgan M. A brief psycho-educational group intervention for postnatal depression. *British Journal of Clinical Psychology*. 2002;41(4):405-9.
255. Hopko DR, Armento ME, Robertson SM, Ryba MM, Carvalho JP, Colman LK, et al. Brief behavioral activation and problem-solving therapy for depressed breast cancer patients: randomized trial. *Journal of consulting and clinical psychology*. 2011;79(6):834-49.
256. Hopko DR, Lejuez CW, LePage JP, Hopko SD, McNeil DW. A brief behavioral activation treatment for depression. A randomized pilot trial within an inpatient psychiatric hospital. *Behavior modification*. 2003;27(4):458-69.
257. Horrell L, Goldsmith KA, Tylee AT, Schmidt UH, Murphy CL, Bonin E-M, et al. One-day cognitive-behavioural therapy self-confidence workshops for people with depression: Randomised controlled trial. *The British Journal of Psychiatry*. 2014;204(3):222-33.
258. Hou Y, Hu P, Zhang Y, Lu Q, Wang D, Yin L, et al. Cognitive behavioral therapy in combination with systemic family therapy improves mild to moderate postpartum depression. *Revista brasileira de psiquiatria (Sao Paulo, Brazil : 1999)*. 2014;36(1):47-52.
259. Hsiao F-H, Lai Y-M, Chen Y-T, Yang T-T, Liao S-C, Ho RT, et al. Efficacy of psychotherapy on diurnal cortisol patterns and suicidal ideation in adjustment disorder with depressed mood. *General hospital psychiatry*. 2014;36(2):214-9.
260. Hsiao FH, Jow GM, Lai YM, Chen YT, Wang KC, Ng SM, et al. The long-term effects of psychotherapy added to pharmacotherapy on morning to evening diurnal cortisol patterns in outpatients with major depression. *Psychotherapy and psychosomatics*. 2011;80(3):166-72.
261. Hsu YC, Wang JJ. Physical, affective, and behavioral effects of group reminiscence on depressed institutionalized elders in Taiwan. *Nursing research*. 2009;58(4):294-9.
262. Huang C-Y, Lai H-L, Chen C-I, Lu Y-C, Li S-C, Wang L-W, et al. Effects of motivational enhancement therapy plus cognitive behaviour therapy on depressive symptoms and health-related quality of life in adults with type II diabetes mellitus: A randomised controlled trial. *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care & Rehabilitation*. 2016;25(5):1275-83.
263. Huber D, Henrich G, Clarkin J, Klug G. Psychoanalytic versus psychodynamic therapy for depression: a three-year follow-up study. *Psychiatry*. 2013;76(2):132-49.

264. Hummel J, Weisbrod C, Boesch L, Himpler K, Hauer K, Hautzinger M, et al. AIDE–Acute Illness and Depression in Elderly Patients. Cognitive Behavioral Group Psychotherapy in Geriatric Patients With Comorbid Depression: A Randomized, Controlled Trial. *Journal of the American Medical Directors Association*. 2017;18(4):341-9.
265. Hunter SB, Watkins KE, Hepner KA, Paddock SM, Ewing BA, Osilla KC, et al. Treating depression and substance use: a randomized controlled trial. *Journal of substance abuse treatment*. 2012;43(2):137-51.
266. Hunter SB, Witkiewitz K, Watkins KE, Paddock SM, Hepner KA. The moderating effects of group cognitive–behavioral therapy for depression among substance users. *Psychology of Addictive Behaviors*. 2012;26(4):906-16.
267. Husain N, Chaudhry N, Fatima B, Husain M, Rizwana A, Chaudhry IB, et al. Antidepressant and group psychosocial treatment for depression: A rater blind exploratory RCT from a low income country. *Behavioural and Cognitive Psychotherapy*. 2014;42(6):693-705.
268. Husain N, Zulqernain F, Carter L-A, Chaudhry I, Fatima B, Kiran T, et al. Treatment of maternal depression in urban slums of Karachi, Pakistan: a randomized controlled trial (RCT) of an integrated maternal psychological and early child development intervention. *Asian journal of psychiatry* [Internet]. 2017; 29:[63-70 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/acps.12164>
269. Hwang WC, Myers HF, Chiu E, Mak E, Butner JE, Fujimoto K, et al. Culturally adapted cognitive-behavioral therapy for Chinese Americans with depression: A randomized controlled trial. *Psychiatric Services*. 2015;66(10):1035-42.
270. Iacoviello BM, Murrough JW, Hoch MM, Huryk KM, Collins KA, Cutter GR, et al. A randomized, controlled pilot trial of the Emotional Faces Memory Task: a digital therapeutic for depression. *npj Digital Medicine*. 2018;1(1).
271. Imamura K, Kawakami N, Furukawa TA, Matsuyama Y, Shimazu A, Umanodan R, et al. Effects of an Internet-based cognitive behavioral therapy (iCBT) program in Manga format on improving subthreshold depressive symptoms among healthy workers: a randomized controlled trial. *PLoS One*. 2014;9(5):e97167.
272. Jacobson NS, Dobson K, Fruzzetti AE, Schmalings KB, Salusky S. Marital therapy as a treatment for depression. *Journal of consulting and clinical psychology*. 1991;59(4):547-57.
273. Jacobson NS, Dobson KS, Truax PA, Addis ME, Koerner K, Gollan JK, et al. A component analysis of cognitive-behavioral treatment for depression. *Journal of consulting and clinical psychology*. 1996;64(2):295-304.
274. Jahangard L, Haghghi M, Bajoghli H, Ahmadpanah M, Ghaleiha A, Zarrabian MK, et al. Training emotional intelligence improves both emotional intelligence and depressive symptoms in inpatients with borderline personality disorder and depression. *International journal of psychiatry in clinical practice*. 2012;16(3):197-204.
275. Jahoda A, Hastings R, Hatton C, Cooper S-A, Dagnan D, Zhang R, et al. Comparison of behavioural activation with guided self-help for treatment of depression in adults with intellectual disabilities: A randomised controlled trial. *The Lancet Psychiatry*. 2017;4(12):909-19.
276. Jahoda A, Hastings R, Hatton C, Cooper SA, McMeekin N, Dagnan D, et al. Behavioural activation versus guided self-help for depression in adults with learning disabilities: the beatit RCT. *Health technology assessment*. 2018;22(53):vii-130.
277. Jakobsen JC, Gluud C, Kongerslev M, Larsen KA, Sorensen P, Winkel P, et al. Third-wave cognitive therapy versus mentalisation-based treatment for major depressive disorder: A randomised clinical trial. *BMJ Open*. 2014;4(8).
278. Jamison C, Scogin F. The outcome of cognitive bibliotherapy with depressed adults. *Journal of consulting and clinical psychology*. 1995;63(4):644-50.
279. Jarrett RB, Schaffer M, McIntire D, Witt-Browder A, Kraft D, Risser RC. Treatment of atypical depression with cognitive therapy or phenelzine: a double-blind, placebo-controlled trial. *Archives of general psychiatry*. 1999;56(5):431-7.
280. Jelinek L, Hauschildt M, Wittekind C, Schneider B, Kriston L, Moritz S. Efficacy of metacognitive training for depression: A randomized controlled trial. *Psychotherapy and psychosomatics* [Internet]. 2016; 85(4):[231-4 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/ppt.12164>
281. Jesse DE, Gaynes BN, Feldhousen EB, Newton ER, Bunch S, Hollon SD. Performance of a Culturally Tailored Cognitive-Behavioral Intervention Integrated in a Public Health Setting to Reduce Risk of Antepartum Depression: A Randomized Controlled Trial. *Journal of Midwifery and Women's Health*. 2015;60(5):578-92.

282. Jiang L, Wang ZZ, Qiu LR, Wan GB, Lin Y, Wei Z. Psychological intervention for postpartum depression. *Journal of Huazhong University of Science and Technology Medical sciences = Hua zhong ke ji da xue xue bao Yi xue Ying De wen ban = Huazhong keji daxue xuebao Yixue Yingdewen ban.* 2014;34(3):437-42.
283. Johansson R, Ekbladh S, Hebert A, Lindström M, Møller S, Pettitt E, et al. Psychodynamic guided self-help for adult depression through the internet: a randomised controlled trial. *PloS one* [Internet]. 2012; 7(5):[e38021 p.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/104/CN-00842104/frame.html>
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3362510/pdf/pone.0038021.pdf>.
284. Johansson R, Sjöberg E, Sjögren M, Johnsson E, Carlbring P, Andersson T, et al. Tailored vs. standardized internet-based cognitive behavior therapy for depression and comorbid symptoms: a randomized controlled trial. *PloS one* [Internet]. 2012; 7(5):[e36905 p.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/787/CN-00852787/frame.html>
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3352859/pdf/pone.0036905.pdf>.
285. Johnson JE, Price AB, Kao JC, Fernandes K, Stout R, Gobin RL, et al. Interpersonal psychotherapy (IPT) for major depression following perinatal loss: a pilot randomized controlled trial. *Archives of Women's Mental Health.* 2016;19(5):845-59.
286. Johnson JE, Zlotnick C. Pilot study of treatment for major depression among women prisoners with substance use disorder. *Journal of psychiatric research.* 2012;46(9):1174-83.
287. Joling KJ, Hout HP, van't Veer-Tazelaar PJ, Horst HE, Cuijpers P, Ven PM, et al. How effective is bibliotherapy for very old adults with subthreshold depression? A randomized controlled trial. *American journal of geriatric psychiatry* [Internet]. 2011; 19(3):[256-65 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/206/CN-00789206/frame.html>
<http://graphics.tx.ovid.com/ovftpdfs/FPDDNCDCGCPFGC00/fs046/ovft/live/gv023/00019442/00019442-201103000-00009.pdf>.
288. Kamga H, McCusker J, Yaffe M, Sewitch M, Sussman T, Strumpf E, et al. Self-care tools to treat depressive symptoms in patients with age-related eye disease: a randomized controlled clinical trial. *Clinical & experimental ophthalmology* [Internet]. 2017; 45(4):[371-8 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/617/CN-01413617/frame.html>
<http://onlinelibrary.wiley.com/store/10.1111/ceo.12890/asset/ceo12890.pdf?v=1&t=je031d5i&s=10a2fa0007b6137283c3a64015e71766f97055bb>.
289. Kanter JW, Santiago-Rivera AL, Santos MM, Nagy G, López M, Hurtado GD, et al. A Randomized Hybrid Efficacy and Effectiveness Trial of Behavioral Activation for Latinos With Depression. *Behavior Therapy.* 2015;46(2):177-92.
290. Karimi H, Dolatshahee B, Momeni K, Khodabakhshi A, Rezaei M, Kamrani A. Effectiveness of integrative and instrumental reminiscence therapies on depression symptoms reduction in institutionalized older adults: An empirical study. *Aging & mental health.* 2010;14(7):881-7.
291. Kay-Lambkin FJ, Baker AL, Kelly B, Lewin TJ. Clinician-assisted computerised versus therapist-delivered treatment for depressive and addictive disorders: a randomised controlled trial. *The Medical journal of Australia.* 2011;195(3):S44-50.
292. Kay-Lambkin FJ, Baker AL, Lewin TJ, Carr VJ. Computer-based psychological treatment for comorbid depression and problematic alcohol and/or cannabis use: a randomized controlled trial of clinical efficacy. *Addiction (Abingdon, England).* 2009;104(3):378-88.
293. Keeley RD, Brody DS, Engel M, Burke BL, Nordstrom K, Morales E, et al. Motivational interviewing improves depression outcome in primary care: A cluster randomized trial. *Journal of Consulting and Clinical Psychology.* 2016;84(11):993-1007.
294. Kelders SM, Bohlmeijer ET, Pots WTM, van Gemert-Pijnen JEW. Comparing human and automated support for depression: Fractional factorial randomized controlled trial. *Behaviour Research and Therapy.* 2015;72:72-80.
295. Keller MB, McCullough JP, Klein DN, Arnow B, Dunner DL, Gelenberg AJ, et al. A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. *The New England journal of medicine.* 2000;342(20):1462-70.
296. Kellett S, Stockton C, Marshall H, Hall J, Jennings C, Degadillo J. Efficacy of narrative reformulation during cognitive analytic therapy for depression: Randomized dismantling trial. *Journal of Affective Disorders.* 2018;239:37-47.
297. Kelly JA, Murphy DA, Bahr GR, Kalichman SC, Morgan MG, Stevenson LY, et al. Outcome of cognitive-behavioral and support group brief therapies for depressed, HIV-infected persons. *The American journal of psychiatry.* 1993;150(11):1679-86.

298. Kennedy SH, Konarski JZ, Segal ZV, Lau MA, Bieling PJ, McIntyre RS, et al. Differences in brain glucose metabolism between responders to CBT and venlafaxine in a 16-week randomized controlled trial. *The American journal of psychiatry*. 2007;164(5):778-88.
299. Kenter RMF, Cuijpers P, Beekman A, van Straten A. Effectiveness of a Web-based guided self-help intervention for outpatients with a depressive disorder: Short-term results from a randomized controlled trial. *Journal of Medical Internet Research*. 2016;18(3).
300. Kessler D, Lewis G, Kaur S, Wiles N, King M, Weich S, et al. Therapist-delivered Internet psychotherapy for depression in primary care: a randomised controlled trial. *Lancet*. 2009;374(9690):628-34.
301. Kim YH, Choi KS, Han K, Kim HW. A psychological intervention programme for patients with breast cancer under chemotherapy and at a high risk of depression: a randomised clinical trial. *Journal of clinical nursing*. 2018;27(3-4):572-81.
302. King M, Sibbald B, Ward E, Bower P, Lloyd M, Gabbay M, et al. Randomised controlled trial of non-directive counselling, cognitive-behaviour therapy and usual general practitioner care in the management of depression as well as mixed anxiety and depression in primary care. *Health technology assessment (Winchester, England)*. 2000;4(19):1-83.
303. Kiosses DN, Arean PA, Teri L, Alexopoulos GS. Home-delivered problem adaptation therapy (PATH) for depressed, cognitively impaired, disabled elders: A preliminary study. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2010;18(11):988-98.
304. Kiosses DN, Ravdin LD, Gross JJ, Raue P, Kotbi N, Alexopoulos GS. Problem adaptation therapy for older adults with major depression and cognitive impairment: a randomized clinical trial. *JAMA psychiatry [Internet]*. 2015; 72(1):[22-30 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/psyc.12007>
305. Kivi M, Eriksson MCM, Hange D, Petersson E-L, Vermark K, Johansson B, et al. Internet-based therapy for mild to moderate depression in Swedish primary care: Short term results from the PRIM-NET randomized controlled trial. *Cognitive Behaviour Therapy*. 2014;43(4):289-98.
306. Klausner EJ, Clarkin JF, Spielman L, Pupo C, Abrams R, Alexopoulos GS. Late-life depression and functional disability: the role of goal-focused group psychotherapy. *International journal of geriatric psychiatry*. 1998;13(10):707-16.
307. Kleiboer A, Donker T, Seekles W, van Straten A, Riper H, Cuijpers P. A randomized controlled trial on the role of support in Internet-based problem solving therapy for depression and anxiety. *Behaviour Research and Therapy*. 2015;72:63-71.
308. Klein J, Berger T, Schröder J, Spith C, Meyer B, Caspar F, et al. Effects of a psychological internet intervention in the treatment of mild to moderate depressive symptoms: Results of the evident study, a randomized controlled trial. *Psychotherapy and psychosomatics [Internet]*. 2016; 85(4):[218-28 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/psyc.12207>
309. Klein MH, Greist JH, Gurman AS, Neimeyer RA, Lesser DP, Bushnell NJ, et al. A comparative outcome study of group psychotherapy vs. exercise treatments for depression. *International Journal of Mental Health*. 1984:148-76.
310. Koenig HG, Pearce MJ, Nelson B, Shaw SF, Robins CJ, Daher NS, et al. Religious vs. conventional cognitive behavioral therapy for major depression in persons with chronic medical illness: A pilot randomized trial. *Journal of nervous and mental disease [Internet]*. 2015; 203(4):[243-51 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/psyc.12007>
311. Konnert C, Dobson K, Stelmach L. The prevention of depression in nursing home residents: a randomized clinical trial of cognitive-behavioral therapy. *Aging & mental health*. 2009;13(2):288-99.
312. Kontunen J, Timonen M, Muotka J, Liukkonen T. Is interpersonal counselling (IPC) sufficient treatment for depression in primary care patients? A pilot study comparing IPC and interpersonal psychotherapy (IPT). *Journal of affective disorders [Internet]*. 2016; 189:[89-93 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/psyc.12207>
313. Korrelboom K, Maarsingh M, Huijbrechts I. Competitive memory training (COMET) for treating low self-esteem in patients with depressive disorders: a randomized clinical trial. *Depression and anxiety*. 2012;29(2):102-10.
314. Korte J, Bohlmeijer ET, Cappeliez P, Smit F, Westerhof GJ. Life review therapy for older adults with moderate depressive symptomatology: a pragmatic randomized controlled trial. *Psychological medicine*

- [Internet]. 2012; 42(6):[1163-73 pp.]. Available from:
<http://onlinelibrary.wiley.com/doi/10.1002/for.1211>
http://journals.cambridge.org/download.php?file=%2FPSM%2FPSM42_06%2FS0033291711002042a.pdf&code=05eaf0d70e2d1087589a28152cc6112a.
315. Koszycki D, Bisscherbe J-C, Blier P, Bradwejn J, Markowitz J. Interpersonal psychotherapy versus brief supportive therapy for depressed infertile women: First pilot randomized controlled trial. *Archives of Women's Mental Health*. 2012;15(3):193-201.
 316. Krampen G. Application of autogenic training before and in addition to integrative psychotherapy of depressive disorders. <ORIGINAL> AUTOGENES TRAINING VOR UND BEGLEITEND ZUR METHODENUBERGREIFENDEN EINZELPSYCHOTHERAPIE BEI DEPRESSIVEN STORUNGEN. *Zeitschrift Fur Klinische Psychologie, Psychiatrie Und Psychotherapie* [Internet]. 1997; 45(2):[214-32 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/for.1211>
 317. Laidlaw K, Davidson K, Toner H, Jackson G, Clark S, Law J, et al. A randomised controlled trial of cognitive behaviour therapy vs treatment as usual in the treatment of mild to moderate late life depression. *International journal of geriatric psychiatry*. 2008;23(8):843-50.
 318. Lam RW, Parikh SV, Ramasubbu R, Michalak EE, Tam EM, Axler A, et al. Effects of combined pharmacotherapy and psychotherapy for improving work functioning in major depressive disorder. *The British journal of psychiatry : the journal of mental science*. 2013;203(5):358-65.
 319. Lamers F, Jonkers CC, Bosma H, Kempen GI, Meijer JA, Penninx BW, et al. A minimal psychological intervention in chronically ill elderly patients with depression: a randomized trial. *Psychotherapy and psychosomatics*. 2010;79(4):217-26.
 320. Lamers SMA, Bohlmeijer ET, Korte J, Westerhof GJ. The efficacy of life-review as online-guided self-help for adults: A randomized trial. *The Journals of Gerontology: Series B: Psychological Sciences and Social Sciences*. 2015;70B(1):24-34.
 321. Landreville P, Bissonnette L. Effects of cognitive bibliotherapy for depressed older adults with a disability. *Clinical Gerontologist*. 1997;17(4):35-55.
 322. Lappalainen P, Granlund A, Siltanen S, Ahonen S, Vitikainen M, Tolvanen A, et al. ACT Internet-based vs face-to-face? A randomized controlled trial of two ways to deliver Acceptance and Commitment Therapy for depressive symptoms: An 18-month follow-up. *Behaviour Research and Therapy*. 2014;61:43-54.
 323. Lappalainen P, Langrial S, Oinas-Kukkonen H, Tolvanen A, Lappalainen R. Web-based acceptance and commitment therapy for depressive symptoms with minimal support: A randomized controlled trial. *Behavior modification*. 2015;39(6):805-34.
 324. Larcombe NA, Wilson PH. An evaluation of cognitive-behaviour therapy for depression in patients with multiple sclerosis. *The British journal of psychiatry : the journal of mental science*. 1984;145:366-71.
 325. Lemma A, Fonagy P. Feasibility study of a psychodynamic online group intervention for depression. *Psychoanalytic Psychology*. 2013;30(3):367-80.
 326. Lemmens GM, Eisler I, Buisse A, Heene E, Demyttenaere K. The effects on mood of adjunctive single-family and multi-family group therapy in the treatment of hospitalized patients with major depression. A 15-month follow-up study. *Psychotherapy and psychosomatics*. 2009;78(2):98-105.
 327. Lemmens LHJM, Arntz A, Peeters F, Hollon SD, Roefs A, Huibers MJH. Clinical effectiveness of cognitive therapy v. interpersonal psychotherapy for depression: Results of a randomized controlled trial. *Psychological Medicine*. 2015;45(10):2095-110.
 328. Lenze S, Potts M. Brief Interpersonal Psychotherapy for depression during pregnancy in a low-income population: a randomized controlled trial. *Journal of affective disorders* [Internet]. 2017; 210:[151-7 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/for.1211>
 329. Lerner D, Adler DA, Rogers WH, Chang H, Greenhill A, Cymerman E, et al. A randomized clinical trial of a telephone depression intervention to reduce employee presenteeism and absenteeism. *Psychiatric Services*. 2015;66(6):570-7.
 330. Lesperance F, Frasere-Smith N, Koszycki D, Laliberte MA, van Zyl LT, Baker B, et al. Effects of citalopram and interpersonal psychotherapy on depression in patients with coronary artery disease: the Canadian Cardiac Randomized Evaluation of Antidepressant and Psychotherapy Efficacy (CREATE) trial. *Jama*. 2007;297(4):367-79.
 331. Leung SS, Lee AM, Chiang VC, Lam SK, Kuen YW, Wong DF. Culturally sensitive, preventive antenatal group cognitive-behavioural therapy for Chinese women with depression. *International journal of nursing practice*. 2013;19 Suppl 1:28-37.

332. Leung SS, Lee AM, Wong DF, Wong CM, Leung KY, Chiang VC, et al. A brief group intervention using a cognitive-behavioural approach to reduce postnatal depressive symptoms: a randomised controlled trial. *Hong Kong medical journal = Xianggang yi xue za zhi*. 2016;22 Suppl 2:S4-8.
333. Levin W, Campbell DR, McGovern KB, Gau JM, Kosty DB, Seeley JR, et al. A computer-assisted depression intervention in primary care. *Psychological medicine*. 2011;41(7):1373-83.
334. Lexis MA, Jansen NW, Huibers MJ, Amelvoort LG, Berkouwer A, Tjin ATG, et al. Prevention of long-term sickness absence and major depression in high-risk employees: a randomised controlled trial. *Occupational and environmental medicine [Internet]*. 2011; 68(6):[400-7 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/941/CN-00788941/frame.html>
<http://oem.bmj.com/content/68/6/400.full.pdf>.
335. Linde JA, Simon GE, Ludman EJ, Ichikawa LE, Operskalski BH, Arterburn D, et al. A randomized controlled trial of behavioral weight loss treatment versus combined weight loss/depression treatment among women with comorbid obesity and depression. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine*. 2011;41(1):119-30.
336. Lindner P, Olsson EL, Johnsson A, Dahlin M, Andersson G, Carlbring P. The impact of telephone versus e-mail therapist guidance on treatment outcomes, therapeutic alliance and treatment engagement in Internet-delivered CBT for depression: A randomised pilot trial. *Internet Interventions [Internet]*. 2014; 1(4):[182-7 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/311/CN-01040311/frame.html>
http://ac.els-cdn.com/S2214782914000256/1-s2.0-S2214782914000256-main.pdf?_tid=54382626-5874-11e5-9b0f-00000aab0f26&acdnat=1441969415_f11f487f885f972a71a25cc88ad507df.
337. Lintvedt OK, Griffiths KM, Sørensen K, Øystvik AR, Wang CE, Eisemann M, et al. Evaluating the effectiveness and efficacy of unguided internet-based self-help intervention for the prevention of depression: a randomized controlled trial. *Clinical psychology & psychotherapy [Internet]*. 2013; 20(1):[10-27 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/770/CN-00905770/frame.html>
<http://onlinelibrary.wiley.com/store/10.1002/cpp.770/asset/cpp770.pdf?v=1&t=i8631rky&s=39315eaf60c11f92ff8e7a4431d2168d919de4fc>
<http://onlinelibrary.wiley.com/store/10.1002/cpp.770/asset/cpp770.pdf?v=1&t=iefj90zk&s=945ca911df17925dd57ac5fe3e679d26a31899f1>.
338. Liu ET-H, Chen W-L, Li Y-H, Wang CH, Mok TJ, Huang HS. Exploring the efficacy of cognitive bibliotherapy and a potential mechanism of change in the treatment of depressive symptoms among the Chinese: A randomized controlled trial. *Cognitive Therapy and Research*. 2009;33(5):449-61.
339. Lloyd-Williams M, Shiels C, Ellis J, Abba K, Gaynor E, Wilson K, et al. Pilot randomised controlled trial of focused narrative intervention for moderate to severe depression in palliative care patients: DISCERN trial. *Palliative medicine*. 2018;32(1):206-15.
340. Lobner M, Pabst A, Stein J, Dorow M, Matschinger H, Luppá M, et al. Computerized cognitive behavior therapy for patients with mild to moderately severe depression in primary care: A pragmatic cluster randomized controlled trial (@ktiv). *Journal of affective disorders*. 2018;238:317-26.
341. Lokman S, Leone S, Sommers-Spijkerman M, Poel A, Smit F, Boon B. Complaint-Directed Mini-Interventions for Depressive Complaints: a Randomized Controlled Trial of Unguided Web-Based Self-Help Interventions. *Journal of medical internet research [Internet]*. 2017; 19(1):[e4 p.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/428/CN-01401428/frame.html>.
342. Lopes RT, Gonçalves MM, Machado PPP, Sinai D, Bento T, Salgado J. Narrative therapy vs. cognitive-behavioral therapy for moderate depression: Empirical evidence from a controlled clinical trial. *Psychotherapy Research*. 2014;24(6):662-74.
343. Losada A, Márquez-González M, Romero-Moreno R, Mausbach BT, López J, Fernández-Fernández V, et al. Cognitive-behavioral therapy (CBT) versus acceptance and commitment therapy (ACT) for dementia family caregivers with significant depressive symptoms: Results of a randomized clinical trial. *Journal of Consulting and Clinical Psychology*. 2015;83(4):760-72.
344. Lovell K, Bower P, Richards D, Barkham Mi, Sibbald B, Roberts C, et al. Developing guided self-help for depression using the medical research council complex interventions framework: A description of the modelling phase and results of an exploratory randomised controlled trial. *BMC psychiatry*. 2008;8.
345. Lundgren J, Dahlstrom Ö, Andersson G, Jaarsma T, Karner Kohlner A, Johansson P. The effect of guided web-based cognitive behavioral therapy on patients with depressive symptoms and heart failure—a randomized controlled trial. *European Journal of Heart Failure*. 2016;18:401.
346. Lustman PJ, Griffith LS, Freedland KE, Kissel SS, Clouse RE. Cognitive behavior therapy for depression in type 2 diabetes mellitus. A randomized, controlled trial. *Annals of internal medicine*. 1998;129(8):613-21.

347. Luty SE, Carter JD, McKenzie JM, Rae AM, Frampton CM, Mulder RT, et al. Randomised controlled trial of interpersonal psychotherapy and cognitive-behavioural therapy for depression. *The British journal of psychiatry : the journal of mental science*. 2007;190:496-502.
348. Luxton DD, Pruitt LD, Wagner A, Smolenski DJ, Jenkins-Guarnieri MA, Gahm G. Home-based telebehavioral health for U.S. military personnel and veterans with depression: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*. 2016;84(11):923-34.
349. Ly KH, Truschel A, Jarl L, Magnusson S, Windahl T, Johansson R, et al. Behavioural activation versus mindfulness-based guided self-help treatment administered through a smartphone application: A randomised controlled trial. *BMJ Open*. 2014;4(1).
350. Lynch D, Tamburrino M, Nagel R, Smith MK. Telephone-based treatment for family practice patients with mild depression. *Psychological reports*. 2004;94(3 Pt 1):785-92.
351. Lynch DJ, Tamburrino MB, Nagel R. Telephone counseling for patients with minor depression: preliminary findings in a family practice setting. *The Journal of family practice*. 1997;44(3):293-8.
352. Lynch TR, Morse JQ, Mendelson T, Robins C. Dialectical behavior therapy for depressed older adults: A randomized pilot study. *American Journal of Geriatric Psychiatry*. 2003;11(1):33-45.
353. Macaskill ND, Macaskill A. Rational-emotive therapy plus pharmacotherapy versus pharmacotherapy alone in the treatment of high cognitive dysfunction depression. *Cognitive Therapy and Research*. 1996;20(6):575-92.
354. MacPherson H, Richmond S, Bland M, Brealey S, Gabe R, Hopton A, et al. Acupuncture and counselling for depression in primary care: a randomised controlled trial. *PLoS medicine*. 2013;10(9):e1001518.
355. Maina G, Forner F, Bogetto F. Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. *Psychotherapy and psychosomatics*. 2005;74(1):43-50.
356. Maina G, Rosso G, Rigardetto S, Chiado Piat S, Bogetto F. No effect of adding brief dynamic therapy to pharmacotherapy in the treatment of obsessive-compulsive disorder with concurrent major depression. *Psychotherapy and psychosomatics*. 2010;79(5):295-302.
357. Maldonado LA. Un modelo de terapia cognitiva desde la perspectiva de la psicología del aprendizaje. *Anuario de psicología/The UB Journal of psychology*. 1984(30):75-96.
358. Maldonado López A. Terapia de conducta y depresión: Un análisis experimental de los modelos conductual y cognitivo. *Revista de Psicología General y Aplicada*. 1982.
359. Malouff JM, Lanyon RI, Schutte NS. Effectiveness of a brief group RET treatment for divorce-related dysphoria. *Journal of rational-emotive and cognitive-behavior therapy*. 1988;6(3):162-71.
360. Manicavasgar V, Parker G, Perich T. Mindfulness-based cognitive therapy vs cognitive behaviour therapy as a treatment for non-melancholic depression. *Journal of Affective Disorders*. 2011;130(1):138-44.
361. Mantani A, Kato T, Furukawa TA, the FI, Horikoshi M, Imai H, et al. Smartphone Cognitive Behavioral Therapy as an Adjunct to Pharmacotherapy for Refractory Depression: Randomized Controlled Trial. *Journal of Medical Internet Research*. 2017;19(11):e373.
362. Markowitz JC, Kocsis JH, Bleiberg KL, Christos PJ, Sacks M. A comparative trial of psychotherapy and pharmacotherapy for "pure" dysthymic patients. *Journal of affective disorders*. 2005;89(1-3):167-75.
363. Markowitz JC, Kocsis JH, Christos P, Bleiberg K, Carlin A. Pilot study of interpersonal psychotherapy versus supportive psychotherapy for dysthymic patients with secondary alcohol abuse or dependence. *The Journal of nervous and mental disease*. 2008;196(6):468-74.
364. Markowitz JC, Kocsis JH, Fishman B, Spielman LA, Jacobsberg LB, Frances AJ, et al. Treatment of depressive symptoms in human immunodeficiency virus-positive patients. *Archives of general psychiatry*. 1998;55(5):452-7.
365. Marshall MB, Zuroff DC, McBride C, Bagby RM. Self-criticism predicts differential response to treatment for major depression. *Journal of clinical psychology*. 2008;64(3):231-44.
366. Martin PR, Aiello R, Gilson K, Meadows G, Milgrom J, Reece J. Cognitive behavior therapy for comorbid migraine and/or tension-type headache and major depressive disorder: An exploratory randomized controlled trial. *Behaviour research and therapy [Internet]*. 2015; 73:[8-18 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/bcpr.12111>
367. Martin SD, Martin E, Rai SS, Richardson MA, Royall R. Brain blood flow changes in depressed patients treated with interpersonal psychotherapy or venlafaxine hydrochloride: preliminary findings. *Archives of general psychiatry*. 2001;58(7):641-8.
368. Matsuzaka C, Wainberg M, Norcini PA, Hoffmann E, Coimbra B, Braga R, et al. Task shifting interpersonal counseling for depression: a pragmatic randomized controlled trial in primary care. *BMC psychiatry [Internet]*. 2017; 17(1) (no pagination). Available from: <http://onlinelibrary.wiley.com/doi/10.1186/s12916-017-0826-6>

- <https://bmcp psychiatry.biomedcentral.com/track/pdf/10.1186/s12888-017-1379-y?site=bmcp psychiatry.biomedcentral.com>.
369. McBride C, Segal Z, Kennedy S, Gemar M. Changes in autobiographical memory specificity following cognitive behavior therapy and pharmacotherapy for major depression. *Psychopathology*. 2007;40(3):147-52.
 370. McClay C-A, Collins K, Matthews L, Haig C, McConnachie A, Morrison J, et al. A community-based pilot randomised controlled study of life skills classes for individuals with low mood and depression. *BMC psychiatry*. 2015;15(1):17.
 371. McCusker J, Cole MG, Yaffe M, Strumpf E, Sewitch M, Sussman T, et al. Adherence to a depression self-care intervention among primary care patients with chronic physical conditions: A randomised controlled trial. *Health Education Journal*. 2016;75(7):767-79.
 372. McCusker J, Cole MG, Yaffe M, Strumpf E, Sewitch M, Sussman T, et al. A randomized trial of a depression self-care toolkit with or without lay telephone coaching for primary care patients with chronic physical conditions. *General hospital psychiatry* [Internet]. 2015; 37(3):[257-65 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/gps.12147>
 373. McGrath CL, Kelley ME, Holtzheimer PE, Dunlop BW, Craighead WE, Franco AR, et al. Toward a neuroimaging treatment selection biomarker for major depressive disorder. *JAMA psychiatry*. 2013;70(8):821-9.
 374. McGrath PJ, Nunes EV, Stewart JW, Goldman D, Agosti V, Ocepek-Welikson K, et al. Imipramine treatment of alcoholics with primary depression: A placebo-controlled clinical trial. *Archives of general psychiatry*. 1996;53(3):232-40.
 375. McIndoo CC, File AA, Preddy T, Clark CG, Hopko DR. Mindfulness-based therapy and behavioral activation: A randomized controlled trial with depressed college students. *Behaviour Research and Therapy*. 2016;77:118-28.
 376. McKee MD, Zayas LH, Fletcher J, Boyd RC, Nam SH. Results of an Intervention to Reduce Perinatal Depression Among Low-Income Minority Women in Community Primary Care. *Journal of Social Service Research*. 2006;32(4):63-81.
 377. McKnight DL, Nelson-Gray RO, Barnhill J. Dexamethasone suppression test and response to cognitive therapy and antidepressant medication. *Behavior Therapy*. 1992;23(1):99-111.
 378. McLean PD, Hakstian AR. Clinical depression: comparative efficacy of outpatient treatments. *Journal of consulting and clinical psychology*. 1979;47(5):818-36.
 379. McNamara K, Horan JJ. Experimental construct validity in the evaluation of cognitive and behavioral treatments for depression. *Journal of Counseling Psychology*. 1986;33(1):23.
 380. Meeks S, Looney SW, Haitsma K, Teri L. BE-ACTIV: a staff-assisted behavioral intervention for depression in nursing homes. *Gerontologist* [Internet]. 2008; 48(1):[105-14 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1532-2207.2007.01405.x>
 381. Meeks S, Van Haitsma K, Schoenbachler B, Looney SW. BE-ACTIV for depression in nursing homes: primary outcomes of a randomized clinical trial. *The journals of gerontology Series B, Psychological sciences and social sciences*. 2015;70(1):13-23.
 382. Menchetti M, Rucci P, Bortolotti B, Bombi A, Scocco P, Kraemer HC, et al. Moderators of remission with interpersonal counselling or drug treatment in primary care patients with depression: randomised controlled trial. *The British journal of psychiatry : the journal of mental science*. 2014;204(2):144-50.
 383. Meyer B, Berger T, Caspar F, Beevers CG, Andersson G, Weiss M. Effectiveness of a novel integrative online treatment for depression (Deprexis): randomized controlled trial. *Journal of medical Internet research*. 2009;11(2):e15.
 384. Meyer B, Bierbrodt J, Schroder J, Berger T, Beevers CG, Weiss M, et al. Effects of an Internet intervention (Deprexis) on severe depression symptoms: Randomized controlled trial. *Internet Interventions* [Internet]. 2015; 2(1):[48-59 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1016/j.invent.2015.01.001>
 385. Michalak J, Schultze M, Heidenreich T, Schramm E. A randomized controlled trial on the efficacy of mindfulness-based cognitive therapy and a group version of cognitive behavioral analysis system of psychotherapy for chronically depressed patients. *Journal of consulting and clinical psychology* [Internet]. 2015; 83(5):[951-63 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/jccp.12147>

386. Milgrom J, Danaher BG, Gemmill AW, Holt C, Holt CJ, Seeley JR, et al. Internet Cognitive Behavioral Therapy for Women With Postnatal Depression: A Randomized Controlled Trial of MumMoodBooster. *Journal of medical Internet research*. 2016;18(3):e54.
387. Milgrom J, Gemmill AW, Ericksen J, Burrows G, Buist A, Reece J. Treatment of postnatal depression with cognitive behavioural therapy, sertraline and combination therapy: a randomised controlled trial. *The Australian and New Zealand journal of psychiatry*. 2015;49(3):236-45.
388. Milgrom J, Holt C, Holt CJ, Ross J, Ericksen J, Gemmill AW. Feasibility study and pilot randomised trial of an antenatal depression treatment with infant follow-up. *Archives of women's mental health*. 2015;18(5):717-30.
389. Milgrom J, Holt CJ, Gemmill AW, Ericksen J, Leigh B, Buist A, et al. Treating postnatal depressive symptoms in primary care: a randomised controlled trial of GP management, with and without adjunctive counselling. *BMC psychiatry*. 2011;11:95.
390. Milgrom J, Negri LM, Gemmill AW, McNeil M, Martin PR. A randomized controlled trial of psychological interventions for postnatal depression. *The British journal of clinical psychology / the British Psychological Society*. 2005;44(Pt 4):529-42.
391. Miller IW, Norman WH, Keitner GI, Bishop SB, Dow MG. Cognitive-behavioral treatment of depressed inpatients. *Behavior Therapy*. 1990;20(1):25-47.
392. Miller L, Weissman M. Interpersonal psychotherapy delivered over the telephone to recurrent depressives. A pilot study. *Depression and anxiety*. 2002;16(3):114-7.
393. Mira A, Breton-Lopez J, Garcia-Palacios A, Quero S, Banos RM, Botella C. An Internet-based program for depressive symptoms using human and automated support: a randomized controlled trial. *Neuropsychiatr Dis Treat*. 2017;13:987-1006.
394. Miranda J, Chung JY, Green BL, Krupnick J, Siddique J, Revicki DA, et al. Treating depression in predominantly low-income young minority women: a randomized controlled trial. *Jama*. 2003;290(1):57-65.
395. Misri S, Kostaras X, Fox D, Kostaras D. The impact of partner support in the treatment of postpartum depression. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2000;45(6):554-8.
396. Misri S, Reebye P, Corral M, Milis L. The use of paroxetine and cognitive-behavioral therapy in postpartum depression and anxiety: a randomized controlled trial. *The Journal of clinical psychiatry*. 2004;65(9):1236-41.
397. Mitchell PH, Veith RC, Becker KJ, Buzaitis A, Cain KC, Fruin M, et al. Brief Psychosocial–Behavioral Intervention With Antidepressant Reduces Poststroke Depression Significantly More Than Usual Care With Antidepressant Living Well With Stroke: Randomized, Controlled Trial. *Stroke; a journal of cerebral circulation*. 2009;40(9):3073-8.
398. Moak DH, Anton RF, Latham PK, Voronin KE, Waid RL, Durazo-Arvizu R. Sertraline and cognitive behavioral therapy for depressed alcoholics: results of a placebo-controlled trial. *Journal of clinical psychopharmacology*. 2003;23(6):553-62.
399. Möbius M, Ferrari GRA, van den Bergh R, Becker ES, Rinck M. Eye-tracking based attention bias modification (ET-ABM) facilitates disengagement from negative stimuli in dysphoric individuals. *Cognitive Therapy and Research*. 2018;42(4):408-20.
400. Mogoșe C, Brăilean A, David D. Can concreteness training alone reduce depressive symptoms? A randomized pilot study using an internet-delivered protocol. *Cognitive therapy and research*. 2013;37(4):704-12.
401. Mohr DC, Boudewyn AC, Goodkin DE, Bostrom A, Epstein L. Comparative outcomes for individual cognitive-behavior therapy, supportive-expressive group psychotherapy, and sertraline for the treatment of depression in multiple sclerosis. *Journal of consulting and clinical psychology*. 2001;69(6):942-9.
402. Mohr DC, Carmody T, Erickson L, Jin L, Leader J. Telephone-administered cognitive behavioral therapy for veterans served by community-based outpatient clinics. *Journal of consulting and clinical psychology*. 2011;79(2):261-5.
403. Mohr DC, Duffecy J, Ho J, Kwasny M, Cai X, Burns MN, et al. A randomized controlled trial evaluating a manualized TeleCoaching protocol for improving adherence to a web-based intervention for the treatment of depression. *PLoS One*. 2013;8(8):e70086.
404. Mohr DC, Hart SL, Julian L, Catledge C, Honos-Webb L, Vella L, et al. Telephone-administered psychotherapy for depression. *Archives of general psychiatry*. 2005;62(9):1007-14.
405. Mohr DC, Ho J, Duffecy J, Reifler D, Sokol L, Burns MN, et al. Effect of telephone-administered vs face-to-face cognitive behavioral therapy on adherence to therapy and depression outcomes among primary care patients: a randomized trial. *Jama*. 2012;307(21):2278-85.
406. Mohr DC, Likosky W, Bertagnolli A, Goodkin DE, Van Der Wende J, Dwyer P, et al. Telephone-administered cognitive-behavioral therapy for the treatment of depressive symptoms in multiple sclerosis. *Journal of consulting and clinical psychology*. 2000;68(2):356-61.

407. Moldovan R, Cobeanu O, David D. Cognitive bibliotherapy for mild depressive symptomatology: randomized clinical trial of efficacy and mechanisms of change. *Clinical psychology & psychotherapy*. 2013;20(6):482-93.
408. Moradveisi L, Huijbers MJ, Renner F, Arasteh M, Arntz A. Behavioural activation v. antidepressant medication for treating depression in Iran: randomised trial. *The British journal of psychiatry : the journal of mental science*. 2013;202(3):204-11.
409. Morgan AJ, Mackinnon AJ, Jorm AF. Behavior change through automated e-mails: Mediation analysis of self-help strategy use for depressive symptoms. *Behaviour research and therapy*. 2013;51(2):57-62.
410. Moritz S, Ahlf-Schumacher J, Hottenrott B, Peter U, Franck S, Schnell T, et al. We cannot change the past, but we can change its meaning. A randomized controlled trial on the effects of self-help imagery rescripting on depression. *Behaviour research and therapy*. 2018;104:74-83.
411. Moritz S, Schilling L, Hauschildt M, Schröder J, Treszl A. A randomized controlled trial of internet-based therapy in depression. *Behaviour research and therapy* [Internet]. 2012; 50(7-8):[513-21 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1469-7610.2012.02522.x>
412. Mossey JM, Knott KA, Higgins M, Talerico K. Effectiveness of a psychosocial intervention, interpersonal counseling, for dysthymic depression in medically ill elderly. *The journals of gerontology Series A, Biological sciences and medical sciences*. 1996;51(4):M172-8.
413. Mukhtar F. Predictors of Group Cognitive Behaviour Therapy outcomes for the treatment of depression in Malaysia. *Asian journal of psychiatry*. 2011;4(2):125-8.
414. Mulcahy R, Reay RE, Wilkinson RB, Owen C. A randomised control trial for the effectiveness of group Interpersonal Psychotherapy for postnatal depression. *Archives of women's mental health*. 2010;13(2):125-39.
415. Murphy GE, Carney RM, Knesevich MA, Wetzel RD, Whitworth P. Cognitive behavior therapy, relaxation training, and tricyclic antidepressant medication in the treatment of depression. *Psychological reports*. 1995;77(2):403-20.
416. Murphy GE, Simons AD, Wetzel RD, Lustman PJ. Cognitive therapy and pharmacotherapy. Singly and together in the treatment of depression. *Archives of general psychiatry*. 1984;41(1):33-41.
417. Myhre MØ, Strömgen B, Arnesen EF, Veland MC. The feasibility of brief behavioural activation treatment for depression in a PICU: A systematic replication. *Journal of Psychiatric Intensive Care*. 2018;14(1):15-23.
418. Mynors-Wallis L, Gath D, Lloyd-Thomas A, Tomlinson D. Randomised controlled trial comparing problem solving treatment with amitriptyline and placebo for major depression in primary care. *Bmj*. 1995;310(6977):441-5.
419. Mynors-Wallis LM, Gath DH, Day A, Baker F. Randomised controlled trial of problem solving treatment, antidepressant medication, and combined treatment for major depression in primary care. *Bmj*. 2000;320(7226):26-30.
420. Naeem F, Gul M, Irfan M, Munshi T, Asif A, Rashid S, et al. Brief Culturally adapted CBT (CaCBT) for depression: A randomized controlled trial from Pakistan. *Journal of affective disorders* [Internet]. 2015; 177:[101-7 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1562-2575.2015.02522.x>
421. Naeem F, Sarhandi I, Gul M, Khalid M, Aslam M, Anbrin A. A multicentre randomised controlled trial of a carer supervised culturally adapted cbt (cacbt) based self-help for depression in pakistan. *Journal of Affective Disorders* [Internet]. 2013; 156:[224-7 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1562-2575.2013.02522.x>
422. Naeem F, Waheed W, Gobbi M, Ayub M, Kingdon D. Preliminary evaluation of culturally sensitive CBT for depression in Pakistan: findings from Developing Culturally-sensitive CBT Project (DCCP). *Behavioural and cognitive psychotherapy*. 2011;39(2):165-73.
423. Nakagawa A, Mitsuda D, Sado M, Abe T, Fujisawa D, Kikuchi T, et al. Effectiveness of supplementary cognitive-behavioral therapy for pharmacotherapy-resistant depression: A randomized controlled trial. *Journal of Clinical Psychiatry*. 2017;78(8):1126-35.
424. Nakimuli-Mpungu E, Wamala K, Okello J, Alderman S, Odokonyero R, Mojtabai R, et al. Group support psychotherapy for depression treatment in people with HIV/AIDS in northern Uganda: a single-centre randomised controlled trial. *The lancet HIV*. 2015;2(5):e190-9.

425. Nasrin F, Rimes K, Reinecke A, Rinck M, Barnhofer T. Effects of Brief Behavioural Activation on Approach and Avoidance Tendencies in Acute Depression: preliminary Findings. *Behavioural and cognitive psychotherapy* [Internet]. 2017; 45(1):[58-72 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/bc.12288>
426. Naylor EV, Antonuccio DO, Litt M, Johnson GE, Spogen DR, Williams R, et al. Bibliotherapy as a treatment for depression in primary care. *Journal of clinical psychology in medical settings*. 2010;17(3):258-71.
427. Neimeyer RA, Feixas G. The role of homework and skill acquisition in the outcome of group cognitive therapy for depression. *Behavior Therapy*. 1990;21(3):281-92.
428. Neugebauer R, Kline J, Markowitz JC, Bleiberg KL, Baxi L, Rosing MA, et al. Pilot randomized controlled trial of interpersonal counseling for subsyndromal depression following miscarriage. *The Journal of clinical psychiatry*. 2006;67(8):1299-304.
429. Newby J, Robins L, Wilhelm K, Smith J, Fletcher T, Gillis I, et al. Web-Based Cognitive Behavior Therapy for Depression in People With Diabetes Mellitus: a Randomized Controlled Trial. *Journal of medical internet research* [Internet]. 2017; 19(5):[e157 p.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/jmir.12888>
430. Newby JM, Lang T, Werner-Seidler A, Holmes E, Moulds ML. Alleviating distressing intrusive memories in depression: A comparison between computerised cognitive bias modification and cognitive behavioural education. *Behaviour research and therapy*. 2014;56:60-7.
431. Nezu AM. Efficacy of a social problem-solving therapy approach for unipolar depression. *Journal of consulting and clinical psychology*. 1986;54(2):196-202.
432. Nezu AM, Perri MG. Social problem-solving therapy for unipolar depression: an initial dismantling investigation. *Journal of consulting and clinical psychology*. 1989;57(3):408-13.
433. Ng SE, Tien A, Thayala JN, Ho RC, Chan MF. The effect of life story review on depression of older community-dwelling Chinese adults in Singapore: a preliminary result. *International journal of geriatric psychiatry*. 2013;28(3):328-30.
434. Ngai FW, Wong P-C, Leung KY, Chau PH, Chung KF. The Effect of Telephone-Based Cognitive-Behavioral Therapy on Postnatal Depression: A Randomized Controlled Trial. *Psychotherapy and psychosomatics* [Internet]. 2015; 84(5):[294-303 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/ppt.12288>
435. Nickel M, Nickel C, Tritt K, Lahmann C, Leiberich P, Loew T, et al. Inpatient treatment of woman with depressive disorders due to stress—does inclusion of the partner in treatment influence outcome? A randomized, controlled, prospective Study. *Wiener Medizinische Wochenschrift*. 2004;154(15-16):378-82.
436. Nobis S, Lehr D, Ebert DD, Baumeister H, Snoek F, Riper H, et al. Efficacy of a web-based intervention with mobile phone support in treating depressive symptoms in adults with type 1 and type 2 diabetes: a randomized controlled trial. *Diabetes care*. 2015;38(5):776-83.
437. Nollett CL, Bray N, Bunce C, Casten RJ, Edwards RT, Hegel MT, et al. Depression in visual impairment trial (Depvit): A randomized clinical trial of depression treatments in people with low vision. *Investigative Ophthalmology and Visual Science*. 2016;57(10):4247-54.
438. Nyström M, Stenling A, Sjöström E, Neely G, Lindner P, Hassmén P, et al. Behavioral activation versus physical activity via the internet: a randomized controlled trial. *Journal of affective disorders* [Internet]. 2017; 215:[85-93 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/ja.12888>
439. O'Hara MW, Stuart S, Gorman LL, Wenzel A. Efficacy of interpersonal psychotherapy for postpartum depression. *Archives of general psychiatry*. 2000;57(11):1039-45.
440. O'Mahen H, Himle JA, Fedock G, Henshaw E, Flynn H. A pilot randomized controlled trial of cognitive behavioral therapy for perinatal depression adapted for women with low incomes. *Depression and anxiety*. 2013;30(7):679-87.
441. O'Mahen HA, Woodford J, McGinley J, Warren FC, Richards DA, Lynch TR, et al. Internet-based behavioral activation--treatment for postnatal depression (Netmums): a randomized controlled trial. *Journal of affective disorders*. 2013;150(3):814-22.
442. O'Moore K A, Newby JM, Andrews G, Hunter DJ, Bennell K, Smith J, et al. Internet Cognitive-Behavioral Therapy for Depression in Older Adults With Knee Osteoarthritis: A Randomized Controlled Trial. *Arthritis care & research*. 2018;70(1):61-70.

443. O'Neil A, Taylor B Fau - Sanderson K, Sanderson K Fau - Cyril S, Cyril S Fau - Chan B, Chan B Fau - Hawkes AL, Hawkes Al Fau - Hare DL, et al. Efficacy and feasibility of a tele-health intervention for acute coronary syndrome patients with depression: results of the "MoodCare" randomized controlled trial. 2014(1532-4796 (Electronic)).
444. Olukolade O, Osinowo H. Efficacy of Cognitive Rehabilitation Therapy on Poststroke Depression among Survivors of First Stroke Attack in Ibadan, Nigeria. *Behavioural neurology* [Internet]. 2017; 2017(no pagination). Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/043/CN-01394043/frame.html>
<http://downloads.hindawi.com/journals/bn/2017/4058124.pdf>.
445. Omidi A, Mohammadkhani P, Mohammadi A, Zargar F. Comparing mindfulness based cognitive therapy and traditional cognitive behavior therapy with treatments as usual on reduction of major depressive disorder symptoms. *Iranian Red Crescent Medical Journal*. 2013;15(2):142-6.
446. Onyechi KCN, Eseadi C, Okere AU, Onuigbo LN, Umoke PCI, Anyaegbunam NJ, et al. Effects of cognitive behavioral coaching on depressive symptoms in a sample of type 2 diabetic inpatients in Nigeria. *Medicine (United States)*. 2016;95(31).
447. Ostacoli L, Carletto S, Cavallo M, Baldomir-Gago P, Di Lorenzo G, Fernandez I, et al. Comparison of eye movement desensitization reprocessing and cognitive behavioral therapy as adjunctive treatments for recurrent depression: The European Depression EMDR Network (EDEN) randomized controlled trial. *Frontiers in Psychology*. 2018;9.
448. Pace TM, Dixon DN. Changes in depressive self-schemata and depressive symptoms following cognitive therapy. *Journal of Counseling Psychology*. 1993;40(3):288.
449. Padfield M. The comparative effects of two counseling approaches on the intensity of depression among rural women of low socioeconomic status. *Journal of Counseling Psychology*. 1976;23(3):209.
450. Pagoto S, Schneider KL, Whited MC, Oleski JL, Merriam P, Appelhans B, et al. Randomized controlled trial of behavioral treatment for comorbid obesity and depression in women: the Be Active Trial. *International journal of obesity (2005)*. 2013;37(11):1427-34.
451. Pardini J, Scogin F, Schriver J, Domino M, Wilson D, LaRocca M. Efficacy and process of cognitive bibliotherapy for the treatment of depression in jail and prison inmates. *Psychological services*. 2014;11(2):141-52.
452. Parker G, Blanch B, Paterson A, Hadzi-Pavlovic D, Sheppard E, Manicavasagar V, et al. The superiority of antidepressant medication to cognitive behavior therapy in melancholic depressed patients: A 12-week single-blind randomized study. *Acta Psychiatrica Scandinavica*. 2013;128(4):271-81.
453. Patel V, Weobong B, Weiss H, Anand A, Bhat B, Katti B, et al. The Healthy Activity Program (HAP), a lay counsellor-delivered brief psychological treatment for severe depression, in primary care in India: a randomised controlled trial. *Lancet (london, england)* [Internet]. 2017; 389(10065):[176-85 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/650/CN-01297650/frame.html>.
454. Pecheur DR, Edwards KJ. A comparison of secular and religious versions of cognitive therapy with depressed Christian college students. *Journal of Psychology and Theology*. 1984.
455. Peden AR, Hall LA, Rayens MK, Beebe LL. Reducing negative thinking and depressive symptoms in college women. *Journal of nursing scholarship : an official publication of Sigma Theta Tau International Honor Society of Nursing / Sigma Theta Tau*. 2000;32(2):145-51.
456. Penckofer SM, Ferrans C, Mumby P, Byrn M, Emanuele MA, Harrison PR, et al. A psychoeducational intervention (SWEEP) for depressed women with diabetes. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine*. 2012;44(2):192-206.
457. Pentecost C, Farrand P, Greaves CJ, Taylor RS, Warren FC, Hillsdon M, et al. Combining behavioural activation with physical activity promotion for adults with depression: Findings of a parallel-group pilot randomised controlled trial (BAcPAc). *Trials* [Internet]. 2015; 16(1). Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/170/CN-01090170/frame.html>
<http://www.trialsjournal.com/content/pdf/s13063-015-0881-0.pdf>.
458. Perini S, Titov N, Andrews G. Clinician-assisted Internet-based treatment is effective for depression: randomized controlled trial. *The Australian and New Zealand journal of psychiatry*. 2009;43(6):571-8.
459. Petersen I, Hanass Hancock J, Bhana A, Govender K. A group-based counselling intervention for depression comorbid with HIV/AIDS using a task shifting approach in South Africa: a randomized controlled pilot study. *Journal of affective disorders*. 2014;158:78-84.
460. Petrak F, Herpertz S, Albus C, Hermanns N, Hiemke C, Hiller W, et al. Cognitive Behavioral Therapy Versus Sertraline in Patients With Depression and Poorly Controlled Diabetes: The Diabetes and Depression (DAD) Study: A Randomized Controlled Multicenter Trial. *Diabetes care*. 2015;38(5):767-75.
461. Phillips R, Schneider J, Molosankwe I, Leese M, Froushani P, Grime P. Randomized controlled trial of computerized cognitive behavioural therapy for depressive symptoms: Effectiveness and costs of a

- workplace intervention. *Psychological Medicine* [Internet]. 2014; 44(4):[741-52 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-3113.2014.05431.x>
462. Pibernik-Okanovic M, Begic D, Ajdukovic D, Andrijasevic N, Metelko Z. Psychoeducation versus treatment as usual in diabetic patients with subthreshold depression: preliminary results of a randomized controlled trial. *Trials*. 2009;10:78.
463. Pibernik-Okanovic M, Hermanns N, Ajdukovic D, Kos J, Prasek M, Sekerija M, et al. Does treatment of subsyndromal depression improve depression-related and diabetes-related outcomes? A randomised controlled comparison of psychoeducation, physical exercise and enhanced treatment as usual. *Trials*. 2015;16:305.
464. Pinheiro RT, Botella L, de Avila Quevedo L, Pinheiro KAT, Jansen K, Osório CM, et al. Maintenance of the effects of cognitive behavioral and relational constructivist psychotherapies in the treatment of women with postpartum depression: A randomized clinical trial. *Journal of Constructivist Psychology*. 2014;27(1):59-68.
465. Pinniger R, Brown RF, Thorsteinsson EB, McKinley P. Argentine tango dance compared to mindfulness meditation and a waiting-list control: a randomised trial for treating depression. *Complementary therapies in medicine*. 2012;20(6):377-84.
466. Polshuck EL, Gamble SA, Bellenger K, Lu N, Tu X, Sorensen S, et al. Randomized controlled trial of interpersonal psychotherapy versus enhanced treatment as usual for women with co-occurring depression and pelvic pain. *Journal of psychosomatic research*. 2014;77(4):264-72.
467. Pot AM, Bohlmeijer ET, Onrust S, Melenhorst A-S, Veerbeek M, De Vries W. The impact of life review on depression in older adults: A randomized controlled trial. *International Psychogeriatrics*. 2010;22(4):572-81.
468. Pots W, Fledderus M, Meulenbeek P, Klooster P, Schreurs K, Bohlmeijer E. Acceptance and commitment therapy as a web-based intervention for depressive symptoms: randomised controlled trial. *The British journal of psychiatry : the journal of mental science* [Internet]. 2016; 208(1):[69-77 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/bjpp.12282>
469. Pots WTM, Meulenbeek PAM, Veehof MM, Klungers J, Bohlmeijer ET. The efficacy of mindfulness-based cognitive therapy as a public mental health intervention for adults with mild to moderate depressive symptomatology: A randomized controlled trial. *PLoS ONE*. 2014;9(10).
470. Power Michael J, Freeman C. A randomized controlled trial of IPT versus CBT in primary care: With some cautionary notes about handling missing values in clinical trials. [References]. *Clinical psychology & psychotherapy* [Internet]. 2012; 19(2):[159-69 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1469-7610.2011.02481.x>
471. Prendergast J, Austin MP. Early childhood nurse-delivered cognitive behavioural counselling for post-natal depression. *Australasian Psychiatry*. 2001;9(3):255-9.
472. Preschl B, Maercker A, Wagner B, Forstmeier S, Banos RM, Alcaniz M, et al. Life-review therapy with computer supplements for depression in the elderly: a randomized controlled trial. *Aging & mental health*. 2012;16(8):964-74.
473. Propst LR, Ostrom R, Watkins P, Dean T, Mashburn D. Comparative efficacy of religious and nonreligious cognitive-behavioral therapy for the treatment of clinical depression in religious individuals. *Journal of consulting and clinical psychology*. 1992;60(1):94-103.
474. Puckering C, McIntosh E, Hickey A, Longford J. Mellow Babies: A group intervention for infants and mothers experiencing postnatal depression. *Counselling Psychology Review*. 2010;25(1):28-38.
475. Pugh NE, Hadjistavropoulos HD, Dirkse D. A Randomised Controlled Trial of Therapist-Assisted, Internet-Delivered Cognitive Behavior Therapy for Women with Maternal Depression. *PLoS One*. 2016;11(3):e0149186.
476. Qiu J, Chen W, Gao X, Xu Y, Tong H, Yang M, et al. A randomized controlled trial of group cognitive behavioral therapy for Chinese breast cancer patients with major depression. *Journal of psychosomatic obstetrics and gynaecology*. 2013;34(2):60-7.
477. Quilty LC, Dozois DJA, Lobo DSS, Ravindran LN, Bagby RM. Cognitive structure and processing during cognitive behavioral therapy vs. pharmacotherapy for depression. *International Journal of Cognitive Therapy*. 2014;7(3):235-50.
478. Rahman A, Malik A, Sikander S, Roberts C, Creed F. Cognitive behaviour therapy-based intervention by community health workers for mothers with depression and their infants in rural Pakistan: a cluster-randomised controlled trial. *Lancet*. 2008;372(9642):902-9.

479. Ransom D, Heckman TG, Anderson T, Garske J, Holroyd K, Basta T. Telephone-delivered, interpersonal psychotherapy for HIV-infected rural persons with depression: a pilot trial. *Psychiatric services* (Washington, DC). 2008;59(8):871-7.
480. Ravindran AV, Anisman H, Merali Z, Charbonneau Y, Telner J, Bialik RJ, et al. Treatment of primary dysthymia with group cognitive therapy and pharmacotherapy: clinical symptoms and functional impairments. *The American journal of psychiatry*. 1999;156(10):1608-17.
481. Rector NA, Cassin SE, Richter MA. Psychological treatment of obsessive-compulsive disorder in patients with major depression: a pilot randomized controlled trial. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2009;54(12):846-51.
482. Rehm LP, Kornblith SJ, O'Hara MW, Lamparski DM, Romano JM, Volkin JI. An evaluation of major components in a self-control therapy program for depression. *Behavior modification*. 1981;5(4):459-89.
483. Reinhardt JP, Horowitz A, Cimarolli VR, Eimicke JP, Teresi JA. Addressing depression in a long-term care setting: A Phase II pilot of problem-solving treatment. *Clinical Therapeutics*. 2014;36(11):1531-7.
484. Reynolds CF, 3rd, Miller MD, Pasternak RE, Frank E, Perel JM, Cornes C, et al. Treatment of bereavement-related major depressive episodes in later life: a controlled study of acute and continuation treatment with nortriptyline and interpersonal psychotherapy. *The American journal of psychiatry*. 1999;156(2):202-8.
485. Reynolds CF, 3rd, Thomas SB, Morse JQ, Anderson SJ, Albert S, Dew MA, et al. Early intervention to preempt major depression among older black and white adults. *Psychiatric services* (Washington, DC). 2014;65(6):765-73.
486. Richards D, Timulak L, O'Brien E, Hayes C, Vigano N, Sharry J, et al. A randomized controlled trial of an internet-delivered treatment: Its potential as a low-intensity community intervention for adults with symptoms of depression. *Behaviour research and therapy* [Internet]. 2015; 75:[20-31 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/bbr.12182>
487. Richards DA, Ekers D, McMillan D, Taylor RS, Byford S, Warren FC, et al. Cost and Outcome of Behavioural Activation versus Cognitive Behavioural Therapy for Depression (COBRA): a randomised, controlled, non-inferiority trial. *Lancet*. 2016;388(10047):871-80.
488. Richards SH, Dickens C, Anderson R, Richards DA, Taylor RS, Ukoumunne OC, et al. Assessing the effectiveness of Enhanced Psychological Care for patients with depressive symptoms attending cardiac rehabilitation compared with treatment as usual (CADENCE): a pilot cluster randomised controlled trial. *Trials*. 2018;19(1).
489. Rief W, Bleichhardt G, Dannehl K, Euteneuer F, Wambach K. Comparing the Efficacy of CBASP with Two Versions of CBT for Depression in a Routine Care Center: a Randomized Clinical Trial. *Psychotherapy and psychosomatics*. 2018;(no pagination).
490. Rieu J, Bui E, Rouch V, Faure K, Birmes P, Schmitt L. Efficacy of ultrabrief cognitive and behavioural therapy performed by psychiatric residents on depressed inpatients. *Psychotherapy and psychosomatics*. 2011;80(6):374-6.
491. Rizvi SJ, Zaretsky A, Schaffer A, Levitt A. Is immediate adjunctive CBT more beneficial than delayed CBT in treating depression?: A Pilot Study. *J Psychiatr Pract*. 2015;21(2):107-13.
492. Rodríguez Vega B, Palao A, Torres G, Benito G, Pérez E, Dieguez M, et al. Combined therapy versus usual care for the treatment of depression in oncologic patients: a randomized controlled trial. *Psycho-oncology* [Internet]. 2011; 20(9):[943-52 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-3113.2011.04611.x>
493. Roepke AM, Jaffee SR, Riffle OM, McGonigal J, Broome R, Maxwell B. Randomized controlled trial of SuperBetter, a smartphone-based/internet-based self-help tool to reduce depressive symptoms. *Games for Health*. 2015;4(3):235-46.
494. Rohan KJ, Mahon JN, Evans M, Ho SY, Meyerhoff J, Postolache TT, et al. Randomized trial of cognitive-behavioral therapy versus light therapy for seasonal affective disorder: Acute outcomes. *American journal of psychiatry* [Internet]. 2015; 172(9 // R01 MH-078982 (NIMH) *National Institute of Mental Health*):[862-9 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/ajp.12400>
495. Rohan KJ, Roecklein KA, Lindsey KT, Johnson LG, Lippy RD, Lacy TJ, et al. A Randomized Controlled Trial of Cognitive-Behavioral Therapy, Light Therapy, and Their Combination for Seasonal Affective Disorder. *Journal of Consulting and Clinical Psychology*. 2007;75(3):489-500.
496. Rohde P, Stice E, Shaw H, Gau JM. Pilot trial of a dissonance-based cognitive-behavioral group depression prevention with college students. *Behaviour Research and Therapy*. 2016;82:21-7.

497. Rohricht F, Papadopoulos N, Priebe S. An exploratory randomized controlled trial of body psychotherapy for patients with chronic depression. *Journal of affective disorders*. 2013;151(1):85-91.
498. Ross M, Scott M. An evaluation of the effectiveness of individual and group cognitive therapy in the treatment of depressed patients in an inner city health centre. *The Journal of the Royal College of General Practitioners*. 1985;35(274):239-42.
499. Rosso G, Martini B, Maina G. Brief dynamic therapy and depression severity: a single-blind, randomized study. *Journal of affective disorders*. 2013;147(1-3):101-6.
500. Rosso I, Killgore W, Olson E, Webb C, Fukunaga R, Auerbach R, et al. Internet-based cognitive behavior therapy for major depressive disorder: a randomized controlled trial. *Depression and anxiety [Internet]*. 2017; 34(3):[236-45 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/da.22701>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5540163/pdf/nihms884093.pdf>
501. Roten Y, Ambresin G, Herrera F, Fassassi S, Fournier N, Preisig M, et al. Efficacy of an adjunctive brief psychodynamic psychotherapy to usual inpatient treatment of depression: results of a randomized controlled trial. *Journal of affective disorders [Internet]*. 2017; 209:[105-13 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/da.22701>
502. Roth D, Bielski R, Jones M. A comparison of self-control therapy and combined self-control therapy and antidepressant medication in the treatment of depression. *Behavior Therapy*. 1982;13(2):133-44.
503. Rovner BW, Casten RJ, Hegel MT, Massof RW, Leiby BE, Ho AC, et al. Low vision depression prevention trial in age-related macular degeneration: a randomized clinical trial. *Ophthalmology*. 2014;121(11):2204-11.
504. Rude SS. Relative benefits of assertion or cognitive self-control treatment for depression as a function of proficiency in each domain. *Journal of consulting and clinical psychology*. 1986;54(3):390-4.
505. Rush AJ, Beck AT, Kovacs M, Hollon S. Comparative efficacy of cognitive therapy and pharmacotherapy in the treatment of depressed outpatients. *Cognitive therapy and research*. 1977;1(1):17-37.
506. Rush AJ, Watkins JT. Group versus individual cognitive therapy: A pilot study. *Cognitive Therapy and Research*. 1981;5(1):95-103.
507. Ruwaard J, Schrieken B, Schrijver M, Broeksteeg J, Dekker J, Vermeulen H, et al. Standardized web-based cognitive behavioural therapy of mild to moderate depression: a randomized controlled trial with a long-term follow-up. *Cognitive behaviour therapy*. 2009;38(4):206-21.
508. Sadler P, McLaren S, Klein B, Harvey J, Jenkins M. Cognitive behavior therapy for older adults with insomnia and depression: a randomized controlled trial in community mental health services. *Sleep*. 2018;41(8) (no pagination).
509. Safren SA, Bedoya CA, O'Cleirigh C, Biello KB, Pinkston MM, Stein MD, et al. Cognitive behavioural therapy for adherence and depression in patients with HIV: a three-arm randomised controlled trial. *The Lancet HIV*. 2016;3(11):e529-e38.
510. Safren SA, Gonzalez JS, Wexler DJ, Psaros C, Delahanty LM, Blashill AJ, et al. A randomized controlled trial of cognitive behavioral therapy for adherence and depression (CBT-AD) in patients with uncontrolled type 2 diabetes. *Diabetes care*. 2014;37(3):625-33.
511. Safren SA, O'Cleirigh C, Tan JY, Raminani SR, Reilly LC, Otto MW, et al. A randomized controlled trial of cognitive behavioral therapy for adherence and depression (CBT-AD) in HIV-infected individuals. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 2009;28(1):1-10.
512. Salkovskis P, Rimes K, Stephenson D, Sacks G, Scott J. A randomized controlled trial of the use of self-help materials in addition to standard general practice treatment of depression compared to standard treatment alone. *Psychological medicine*. 2006;36(03):325-33.
513. Salminen JK, Karlsson H, Hietala J, Kajander J, Aalto S, Markkula J, et al. Short-term psychodynamic psychotherapy and fluoxetine in major depressive disorder: a randomized comparative study. *Psychotherapy and psychosomatics*. 2008;77(6):351-7.
514. Saloheimo HP, Markowitz J, Saloheimo TH, Laitinen JJ, Sundell J, Huttunen MO, et al. Psychotherapy effectiveness for major depression: a randomized trial in a Finnish community. *BMC psychiatry*. 2016;16:131.
515. Sandoval L, Buckey J, Ainslie R, Tombari M, Stone W, Hegel M. Randomized Controlled Trial of a Computerized Interactive Media-Based Problem Solving Treatment for Depression. *Behavior therapy [Internet]*. 2017; 48(3):[413-25 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/da.22701>
https://ac.els-cdn.com/S0005789416300090/1-s2.0-S0005789416300090-main.pdf?_tid=eb56b75e-b991-11e7-8a7c-00000aacb361&acdnat=1508942383_e5bdab491349fc6fda311cea085e73e3

516. Satre DD, Delucchi K, Lichtmacher J, Sterling SA, Weisner C. Motivational interviewing to reduce hazardous drinking and drug use among depression patients. *Journal of substance abuse treatment*. 2013;44(3):323-9.
517. Savard J, Simard S, Giguere I, Ivers H, Morin CM, Maunsell E, et al. Randomized clinical trial on cognitive therapy for depression in women with metastatic breast cancer: psychological and immunological effects. *Palliative & supportive care*. 2006;4(3):219-37.
518. Schaub A, Goldmann U, Mueser TK, Goerigk S, Hautzinger M, Roth E, et al. Efficacy of extended clinical management, group CBT, and group plus individual CBT for major depression: Results of a two-year follow-up study. *Journal of affective disorders*. 2018;238:570-8.
519. Schiffer RB, Wineman NM. Antidepressant pharmacotherapy of depression associated with multiple sclerosis. *The American journal of psychiatry*. 1990;147(11):1493-7.
520. Schmidt MM, Miller WR. Amount of therapist contact and outcome in a multidimensional depression treatment program. *Acta Psychiatrica Scandinavica*. 1983;67(5):319-32.
521. Schmitz JM, Averill P, Stotts AL, Moeller FG, Rhoades HM, Grabowski J. Fluoxetine treatment of cocaine-dependent patients with major depressive disorder. *Drug and alcohol dependence*. 2001;63(3):207-14.
522. Schramm E, Calker D, Dykieriek P, Lieb K, Kech S, Zobel I, et al. An intensive treatment program of interpersonal psychotherapy plus pharmacotherapy for depressed inpatients: acute and long-term results. *American journal of psychiatry* [Internet]. 2007; 164(5):[768-77 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1399-8619.2007.01338.x>
523. Schramm E, Kriston L, Zobel I, Bailer J, Wambach K, Backenstrass M, et al. Effect of Disorder-Specific vs Nonspecific Psychotherapy for Chronic Depression: a Randomized Clinical Trial. *JAMA psychiatry* [Internet]. 2017; 74(3):[233-42 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1001/jamapsychiatry.2017.0160>
https://jamanetwork.com/journals/jamapsychiatry/articlepdf/2600225/jamapsychiatry_Schramm_2017_oi_160099.pdf
524. Schramm E, Zobel I, Dykieriek P, Kech S, Brakemeier EL, Kulz A, et al. Cognitive behavioral analysis system of psychotherapy versus interpersonal psychotherapy for early-onset chronic depression: a randomized pilot study. *Journal of affective disorders*. 2011;129(1-3):109-16.
525. Schramm E, Zobel I, Schoepf D, Fangmeier T, Schnell K, Walter H, et al. Cognitive Behavioral Analysis System of Psychotherapy versus Escitalopram in Chronic Major Depression. *Psychotherapy and psychosomatics*. 2015;84(4):227-40.
526. Schroder J, Bruckner K, Fischer A, Lindenau M, Kother U, Vettorazzi E, et al. Efficacy of a psychological online intervention for depression in people with epilepsy: a randomized controlled trial. *Epilepsia*. 2014;55(12):2069-76.
527. Schroevers MJ, Tovote KA, Snippe E, Fleer J. Group and individual mindfulness-based cognitive therapy (MBCT) are both effective: A pilot randomized controlled trial in depressed people with a somatic disease. *Mindfulness*. 2016;7(6):1339-46.
528. Schulberg HC, Block MR, Madonia MJ, Scott CP, Rodriguez E, Imber SD, et al. Treating major depression in primary care practice. Eight-month clinical outcomes. *Archives of general psychiatry*. 1996;53(10):913-9.
529. Schuster R, Leitner I, Carlbring P, Laireiter A-R. Exploring blended group interventions for depression: randomised controlled feasibility study of a blended computer- and multimedia-supported psychoeducational group intervention for adults with depressive symptoms. *Internet interventions* [Internet]. 2017; 8:[63-71 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/inl.12133>
530. Scogin F, Hamblin D, Beutler L. Bibliotherapy for depressed older adults: A self-help alternative. *The Gerontologist*. 1987;27(3):383-7.
531. Scogin F, Jamison C, Gochneaur K. Comparative efficacy of cognitive and behavioral bibliotherapy for mildly and moderately depressed older adults. *Journal of consulting and clinical psychology*. 1989;57(3):403-7.
532. Scogin F, Lichstein K, DiNapoli EA, Woosley J, Thomas SJ, Larocca MA, et al. Effects of integrated telehealth-delivered cognitive-behavioral therapy for depression and insomnia in rural older adults. *Journal of psychotherapy integration*. 2018;28(3):292-309.
533. Scogin FR, Moss K, Harris GM, Presnell AH. Treatment of depressive symptoms in diverse, rural, and vulnerable older adults. *International journal of geriatric psychiatry*. 2014;29(3):310-6.
534. Scott AI, Freeman CP. Edinburgh primary care depression study: treatment outcome, patient satisfaction, and cost after 16 weeks. *Bmj*. 1992;304(6831):883-7.

535. Scott C, Tacchi MJ, Jones R, Scott J. Acute and one-year outcome of a randomised controlled trial of brief cognitive therapy for major depressive disorder in primary care. *The British journal of psychiatry : the journal of mental science*. 1997;171:131-4.
536. Scott MJ, Stradling SG. Group cognitive therapy for depression produces clinically significant reliable change in community-based settings. *Behavioural Psychotherapy*. 1990;18(01):1-19.
537. Segre LS, Brock RL, O'Hara MW. Depression treatment for impoverished mothers by point-of-care providers: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*. 2015;83(2):314-24.
538. Selmi PM, Klein MH, Greist JH, Sorrell SP, Erdman HP. Computer-administered cognitive-behavioral therapy for depression. *The American journal of psychiatry*. 1990;147(1):51-6.
539. Semkovska M, Lambe S, Lonergain DO, McLoughlin DM. Neurocognitive Remediation Therapy for Depression: A Feasibility Study and Randomized Controlled Pilot Protocol Testing. *The Journal of nervous and mental disease*. 2015;203(8):609-16.
540. Serfaty MA, Haworth D, Blanchard M, Buszewicz M, Murad S, King M. Clinical effectiveness of individual cognitive behavioral therapy for depressed older people in primary care: a randomized controlled trial. *Archives of general psychiatry*. 2009;66(12):1332-40.
541. Serrano JP, Latorre JM, Gatz M, Montanes J. Life review therapy using autobiographical retrieval practice for older adults with depressive symptomatology. *Psychology and aging*. 2004;19(2):270-7.
542. Serrano Selva JP, Latorre Postigo JM, Ros Segura L, Navarro Bravo B, Aguilar Corcoles MJ, Nieto Lopez M, et al. Life review therapy using autobiographical retrieval practice for older adults with clinical depression. *Psicothema*. 2012;24(2):224-9.
543. Shah A, Morthland M, Scogin F, Presnell A, DiNapoli EA, DeCoster J, et al. Audio and Computer Cognitive Behavioral Therapy for Depressive Symptoms in Older Adults: a Pilot Randomized Controlled Trial. *Behavior therapy*. 2018;49(6):904-16.
544. Shamsaei F, Rahimi A, Zarabian MK, Sedehi M. Efficacy of pharmacotherapy and cognitive therapy, alone and in combination in major depressive disorder. *Hong Kong Journal of Psychiatry*. 2008;18(2):76-80.
545. Shapiro DA, Barkham M, Hardy GE, Morrison LA. The Second Sheffield Psychotherapy Project: rationale, design and preliminary outcome data. *The British journal of medical psychology*. 1990;63 (Pt 2):97-108.
546. Sharp DJ, Chew-Graham C, Tylee A, Lewis G, Howard L, Anderson I, et al. A pragmatic randomised controlled trial to compare antidepressants with a community-based psychosocial intervention for the treatment of women with postnatal depression: the RESPOND trial. *Health technology assessment (Winchester, England)*. 2010;14(43):iii-iv, ix-xi, 1-153.
547. Shaw BF. Comparison of cognitive therapy and behavior therapy in the treatment of depression. *Journal of Consulting and Clinical Psychology*. 1977;45(4):543.
548. Sheeber L, Feil E, Seeley J, Leve C, Gau J, Davis B, et al. Mom-net: evaluation of an internet-facilitated cognitive behavioral intervention for low-income depressed mothers. *Journal of consulting and clinical psychology [Internet]*. 2017; 85(4):[355-66 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/9781119281111.ch43>
549. Sheeber LB, Seeley JR, Feil EG, Davis B, Sorensen E, Kosty DB, et al. Development and pilot evaluation of an Internet-facilitated cognitive-behavioral intervention for maternal depression. *Journal of consulting and clinical psychology*. 2012;80(5):739-49.
550. Silverstone PH, Rittenbach K, Suen VYM, Moretzsohn A, Cribben I, Bercov M, et al. Depression outcomes in adults attending family practice were not improved by screening, stepped-care, or online CBT during a 12-week study when compared to controls in a randomized trial. *Frontiers in Psychiatry*. 2017;8(MAR).
551. Simoni JM, Wiebe JS, Saucedo JA, Huh D, Sanchez G, Longoria V, et al. A preliminary RCT of CBT-AD for adherence and depression among HIV-positive Latinos on the U.S.-Mexico border: the Nuevo Dia study. *AIDS and behavior*. 2013;17(8):2816-29.
552. Simpson S, Corney R, Beecham J. A randomized controlled trial to evaluate the effectiveness and cost-effectiveness of psychodynamic counselling for general practice patients with chronic depression. *Psychological Medicine*. 2003;33(2):229-39.
553. Simson U, Nawarotzky U, Friese G, Porck W, Schottenfeld-Naor Y, Hahn S, et al. Psychotherapy intervention to reduce depressive symptoms in patients with diabetic foot syndrome. *Diabetic medicine : a journal of the British Diabetic Association*. 2008;25(2):206-12.
554. Singh NA, Clements KM, Fiatarone MA. A randomized controlled trial of progressive resistance training in depressed elders. *Journals of gerontology Series A, Biological sciences and medical sciences [Internet]*. 1997; 52(1):[M27-35 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/9781119281111.ch43>

555. Sinniah A, Oei T, Maniam T, Subramaniam P. Positive effects of Individual Cognitive Behavior Therapy for patients with unipolar mood disorders with suicidal ideation in Malaysia: a randomised controlled trial. *Psychiatry research* [Internet]. 2017; 254:[179-89 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1016/j.psychres.2017.03.031>
556. Sirey JA, Bruce ML, Alexopoulos GS. The Treatment Initiation Program: an intervention to improve depression outcomes in older adults. *The American journal of psychiatry*. 2005;162(1):184-6.
557. Sloane R, Staples F, Schneider L. Interpersonal therapy versus nortriptyline for depression in the elderly. *Clinical and pharmacological studies in psychiatric disorders*. 1985:344-6.
558. Smit A, Kluiters H, Conradi HJ, Meer K, Tiemens BG, Jenner JA, et al. Short-term effects of enhanced treatment for depression in primary care: results from a randomized controlled trial. *Psychological medicine* [Internet]. 2006; 36(1):[15-26 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-3113.2005.03584.x>
http://journals.cambridge.org/download.php?file=%2FPSM%2FPSM36_01%2FS0033291705006318a.pdf&code=61a29f44e5d071d2ec9cdaa4de02bcf5
559. Smith H, Dillon K, Cogle J. Modification of Hostile Interpretation Bias in Depression: a Randomized Controlled Trial. *Behavior therapy* [Internet]. 2017; (no pagination). Available from: <http://onlinelibrary.wiley.com/doi/10.1002/behb.1238>
https://ac.els-cdn.com/S0005789417300886/1-s2.0-S0005789417300886-main.pdf?_tid=bcda636c-b991-11e7-bff8-0000aacb35e&acdnat=1508942305_31b0e8a7d9157cfc000b8335d8dde873
560. Smith HL, Dillon KH, Cogle JR. Modification of Hostile Interpretation Bias in Depression: a Randomized Controlled Trial. *Behavior therapy*. 2018;49(2):198-211.
561. Smith J, Newby JM, Burston N, Murphy MJ, Michael S, Mackenzie A, et al. Help from home for depression: A randomised controlled trial comparing internet-delivered cognitive behaviour therapy with bibliotherapy for depression. *Internet Interventions*. 2017;9:25-37.
562. Snarski M, Scogin F, DiNapoli E, Presnell A, McAlpine J, Marcinak J. The effects of behavioral activation therapy with inpatient geriatric psychiatry patients. *Behavior therapy*. 2011;42(1):100-8.
563. Soares MC, Mondin TC, Silva G, Barbosa LP, Molina ML, Jansen K, et al. Comparison of Clinical Significance of Cognitive-Behavioral Therapy and Psychodynamic Therapy for Major Depressive Disorder: a Randomized Clinical Trial. *Journal of nervous and mental disease*. 2018;206(9):686-93.
564. Songprakun W, McCann TV. Evaluation of a cognitive behavioural self-help manual for reducing depression: a randomized controlled trial. *Journal of psychiatric and mental health nursing*. 2012;19(7):647-53.
565. Souza L, Salum G, Mosqueiro B, Caldieraro M, Guerra T, Fleck M. Interpersonal psychotherapy as add-on for treatment-resistant depression: A pragmatic randomized controlled trial. *Journal of affective disorders* [Internet]. 2016; 193:[373-80 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1016/j.jad.2016.03.031>
http://ac.els-cdn.com/S0165032716000045/1-s2.0-S0165032716000045-main.pdf?_tid=baa9606a-d9a0-11e6-9479-0000aacb35e&acdnat=1484319683_01c7defe1d63f700d8f999ab09edb538
566. Spek V, Nyklicek I, Smits N, Cuijpers P, Riper H, Keyzer J, et al. Internet-based cognitive behavioural therapy for subthreshold depression in people over 50 years old: a randomized controlled clinical trial. *Psychological medicine*. 2007;37(12):1797-806.
567. Spinelli MG, Endicott J. Controlled clinical trial of interpersonal psychotherapy versus parenting education program for depressed pregnant women. *The American journal of psychiatry*. 2003;160(3):555-62.
568. Spinelli MG, Endicott J, Leon AC, Goetz RR, Kalish RB, Brustman LE, et al. A controlled clinical treatment trial of interpersonal psychotherapy for depressed pregnant women at 3 New York City sites. *The Journal of clinical psychiatry*. 2013;74(4):393-9.
569. Sreevani R, Reddemma K, Chan CL, Leung PP, Wong V, Chan CH. Effectiveness of integrated body-mind-spirit group intervention on the well-being of Indian patients with depression: a pilot study. *The journal of nursing research : JNR*. 2013;21(3):179-86.
570. Stiles-Shields C, Montague E, Kwasny MJ, Mohr DC. Behavioral and cognitive intervention strategies delivered via coached apps for depression: Pilot trial. *Psychological services*. 2018.
571. Strauman TJ, Vieth AZ, Merrill KA, Kolden GG, Woods TE, Klein MH, et al. Self-system therapy as an intervention for self-regulatory dysfunction in depression: a randomized comparison with cognitive therapy. *Journal of consulting and clinical psychology*. 2006;74(2):367-76.
572. Strauss C, Hayward M, Chadwick P. Group person-based cognitive therapy for chronic depression: a pilot randomized controlled trial. *The British journal of clinical psychology / the British Psychological Society*. 2012;51(3):345-50.

573. Stravynski A, Verreault R, Gaudette G, Langlois R, Gagnier S, Larose M. The treatment of depression with group behavioural-cognitive therapy and imipramine. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 1994;39(7):387-90.
574. Strong V, Waters R, Hibberd C, Murray G, Wall L, Walker J, et al. Management of depression for people with cancer (SMaRT oncology 1): a randomised trial. *Lancet*. 2008;372(9632):40-8.
575. Sugg HVR, Richards DA, Frost J. Morita Therapy for depression (Morita Trial): a pilot randomised controlled trial. *BMJ open*. 2018;8(8).
576. Swartz HA, Cyranowski JM, Cheng Y, Zuckoff A, Brent DA, Markowitz JC, et al. Brief psychotherapy for maternal depression: Impact on mothers and children. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2016;55(6):495-503.
577. Swartz HA, Frank E, Zuckoff A, Cyranowski JM, Houck PR, Cheng Y, et al. Brief interpersonal psychotherapy for depressed mothers whose children are receiving psychiatric treatment. *The American journal of psychiatry*. 2008;165(9):1155-62.
578. Takagaki K, Okamoto Y, Jinnin R, Mori A, Nishiyama Y, Yamamura T, et al. Behavioral activation for late adolescents with subthreshold depression: a randomized controlled trial. *European Child and Adolescent Psychiatry*. 2016;25(11):1171-82.
579. Talbot NL, Chaudron LH, Ward EA, Duberstein PR, Conwell Y, O'Hara MW, et al. A randomized effectiveness trial of interpersonal psychotherapy for depressed women with sexual abuse histories. *Psychiatric services (Washington, DC)*. 2011;62(4):374-80.
580. Targ EF, Karasic DH, Diefenbach PN, Anderson DA, Bystritsky A, Fawzy FI. Structured group therapy and fluoxetine to treat depression in HIV-positive persons. *Psychosomatics*. 1994;35(2):132-7.
581. Taylor BL, Strauss C, Cavanagh K, Jones F. The effectiveness of self-help mindfulness-based cognitive therapy in a student sample: A randomised controlled trial. *Behaviour research and therapy*. 2014;63:63-9.
582. Taylor CB, Conrad A, Wilhelm FH, Strachowski D, Khaylis A, Neri E, et al. Does improving mood in depressed patients alter factors that may affect cardiovascular disease risk? *Journal of psychiatric research*. 2009;43(16):1246-52.
583. Taylor FG, Marshall WL. Experimental analysis of a cognitive-behavioral therapy for depression. *Cognitive Therapy and Research*. 1977;1(1):59-72.
584. Teasdale JD, Fennell MJ, Hibbert GA, Amies PL. Cognitive therapy for major depressive disorder in primary care. *The British journal of psychiatry : the journal of mental science*. 1984;144:400-6.
585. Teichman Y, Bar-el Z, Shor H, Sirota P, Elizur A. A comparison of two modalities of cognitive therapy (individual and marital) in treating depression. *Psychiatry*. 1995;58(2):136-48.
586. Teismann T, Dymel W, Schulte D, Willutzki U. Ressourcenorientierte Akutbehandlung unipolarer Depressionen: Eine randomisierte kontrollierte Psychotherapiestudie. = Resource-focused treatment for unipolar depression: A randomized controlled psychotherapy study. PPM: Psychotherapie Psychosomatik Medizinische Psychologie. 2011;61(7):295-302.
587. Teri L, Lewinsohn PM. Individual and group treatment of unipolar depression: comparison of treatment outcome and identification of predictors of successful treatment outcome. *Behavior therapy*. 1986;17(3):215-28.
588. Teri L, Logsdon RG, Uomoto J, McCurry SM. Behavioral treatment of depression in dementia patients: a controlled clinical trial. *The journals of gerontology Series B, Psychological sciences and social sciences*. 1997;52(4):P159-66.
589. Thase ME, Wright JH, Eells TD, Barrett MS, Wisniewski SR, Balasubramani GK, et al. Improving the Efficiency of Psychotherapy for Depression: computer-Assisted Versus Standard CBT. *American journal of psychiatry*. 2018;175(3):242-50.
590. Thitipitchayanant K, Somrongthong R, Kumar R, Kanchanakharn N. Effectiveness of self-empowerment-affirmation-relaxation(Self-EAR) program for postpartum blues mothers: A randomized controlled trial. *Pakistan Journal of Medical Sciences*. 2018;34(6):1488-93.
591. Thompson LW, Coon DW, Gallagher-Thompson D, Sommer BR, Koin D. Comparison of desipramine and cognitive/behavioral therapy in the treatment of elderly outpatients with mild-to-moderate depression. *The American journal of geriatric psychiatry*. 2001;9(3):225-40.
592. Thompson LW, Gallagher D, Breckenridge JS. Comparative effectiveness of psychotherapies for depressed elders. *Journal of consulting and clinical psychology*. 1987;55(3):385.
593. Thyme KE, Sundin EC, Stahlberg G, Lindstrom B, Eklof H, Wiberg B. The outcome of short-term psychodynamic art therapy compared to short-term psychodynamic verbal therapy for depressed women. *Psychoanalytic Psychotherapy*. 2007;21(3):250-64.
594. Titov N, Andrews G, Davies M, McIntyre K, Robinson E, Solley K. Internet treatment for depression: a randomized controlled trial comparing clinician vs. technician assistance. *PLoS One*. 2010;5(6):e10939.

595. Titov N, Dear BF, Ali S, Zou JB, Lorian CN, Johnston L, et al. Clinical and cost-effectiveness of therapist-guided internet-delivered cognitive behavior therapy for older adults with symptoms of depression: a randomized controlled trial. *Behavior therapy*. 2015;46(2):193-205.
596. Titov N, Dear BF, Staples LG, Terides MD, Karin E, Sheehan J, et al. Disorder-specific versus transdiagnostic and clinician-guided versus self-guided treatment for major depressive disorder and comorbid anxiety disorders: A randomized controlled trial. *Journal of anxiety disorders*. 2015;35:88-102.
597. Tobin K, Davey-Rothwell MA, Nonyane BAS, Knowlton A, Wissow L, Latkin CA. RCT of an integrated CBT-HIV intervention on depressive symptoms and HIV risk. *PLoS One*. 2017;12(12):e0187180.
598. Tomasino K, Lattie E, Ho J, Palac H, Kaiser S, Mohr D. Harnessing Peer Support in an Online Intervention for Older Adults with Depression. *American journal of geriatric psychiatry [Internet]*. 2017; 25(10):[1109-19 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/ajgp.12345>.
599. Torkan H, Blackwell SE, Holmes EA, Kalantari M, Neshat-Doost HT, Maroufi M, et al. Positive imagery cognitive bias modification in treatment-seeking patients with major depression in Iran: A pilot study. *Cognitive Therapy and Research*. 2014;38(2):132-45.
600. Tovote KA, Fleer J, Snippe E, Peeters A, Emmelkamp PMG, Sanderman R, et al. Individual mindfulness-based cognitive therapy and cognitive behavior therapy for treating depressive symptoms in patients with diabetes: Results of a randomized controlled trial. *Diabetes care [Internet]*. 2014; 37(9):[2427-34 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.2337/131122> <http://care.diabetesjournals.org/content/37/9/2427.full.pdf>.
601. Town J, Abbas A, Stride C, Bernier D. A randomised controlled trial of Intensive Short-Term Dynamic Psychotherapy for treatment resistant depression: the Halifax Depression Study. *Journal of affective disorders [Internet]*. 2017; 214:[15-25 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/ajgp.12345>.
602. Trapp W, Engel S, Hajak G, Lautenbacher S, Gallhofer B. Cognitive remediation for depressed inpatients: Results of a pilot randomized controlled trial. *The Australian and New Zealand journal of psychiatry [Internet]*. 2016; 50(1):[46-55 pp.]. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/ajgp.12345>.
603. Travers C. Increasing enjoyable activities to treat depression in nursing home residents with dementia: A pilot study. *Dementia (London, England)*. 2017;16(2):204-18.
604. Trebo E, Holzner B, Pircher M, Prunnelechner R, Gunther V. The effects of a computer assisted cognitive training on neuropsychological parameters, mood and dysfunctional cognitions in depressive patients. *Neuropsychiatrie*. 2007;21(3):207-15.
605. Tsai YF, Wong TK, Tsai HH, Ku YC. Self-worth therapy for depressive symptoms in older nursing home residents. *Journal of advanced nursing*. 2008;64(5):488-94.
606. Tulbure BT, Andersson G, Salagean N, Pearce M, Koenig HG. Religious versus Conventional Internet-based Cognitive Behavioral Therapy for Depression. *Journal of religion and health*. 2018;57(5):1634-48.
607. Turner A, Hambridge J, Baker A, Bowman J, McElduff P. Randomised controlled trial of group cognitive behaviour therapy versus brief intervention for depression in cardiac patients. *The Australian and New Zealand journal of psychiatry*. 2013;47(3):235-43.
608. Turner RW, Ward MF, Turner DJ. Behavioral treatment for depression: An evaluation of therapeutic components. *Journal of clinical psychology*. 1979;35(1):166-75.
609. Tyson GM, Range LM. Gestalt dialogues as a treatment for mild depression: time works just as well. *Journal of clinical psychology*. 1987;43(2):227-31.
610. Unlu Ince B, Cuijpers P, Hof E, Ballegooijen W, Christensen H, Riper H. Internet-based, culturally sensitive, problem-solving therapy for Turkish migrants with depression: randomized controlled trial. *Journal of Medical Internet Research [Internet]*. 2013; 15(10):[e227 p.]. Available from: <http://onlinelibrary.wiley.com/doi/10.2196/jmir.2013.2345>.
611. van Bastelaar KM, Pouwer F, Cuijpers P, Riper H, Snoek FJ. Web-based depression treatment for type 1 and type 2 diabetic patients: a randomized, controlled trial. *Diabetes care*. 2011;34(2):320-5.
612. van Schaik A, van Marwijk H, Adèr H, van Dyck R, de Haan M, Penninx B, et al. Interpersonal Psychotherapy for Elderly Patients in Primary Care. *The American Journal of Geriatric Psychiatry*. 2006;14(9):777-86.
613. Vázquez FL, Torres A, Blanco V, Díaz O, Otero P, Hermida E. Comparison of relaxation training with a cognitive-behavioural intervention for indicated prevention of depression in university students: A randomized controlled trial. *Journal of Psychiatric Research*. 2012;46(11):1456-63.
614. Vázquez FL, Torres Á, Otero P, Blanco V, Díaz O, Estévez LE. Analysis of the components of a cognitive-behavioral intervention administered via conference call for preventing depression among non-professional caregivers: A pilot study. *Aging & mental health*. 2017;21(9):938-46.

615. Vazquez Gonzalez FL, Otero Otero P, Torres Iglesias A, Hermida García E, Blanco Seoane V, Díaz Fernández O. A brief problem-solving indicated-prevention intervention for prevention of depression in nonprofessional caregivers. *Psicothema* [Internet]. 2013; 25(1):[87-92 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/513/CN-00959513/frame.html>.
616. Verduyn C, Barrowclough C, Roberts J, Tarrier N, Harrington R. Maternal depression and child behaviour problems Randomised placebo-controlled trial of a cognitive-behavioural group intervention. *British Journal of Psychiatry*. 2003;183(OCT.):342-8.
617. Vernmark K, Lenndin J, Björrehed J, Carlsson M, Karlsson J, Oberg J, et al. Internet administered guided self-help versus individualized e-mail therapy: A randomized trial of two versions of CBT for major depression. *Behaviour research and therapy* [Internet]. 2010; 48(5):[368-76 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/971/CN-00749971/frame.html>
http://ac.els-cdn.com/S0005796710000069/1-s2.0-S0005796710000069-main.pdf?_tid=2dc862c8-5ade-11e5-92d6-00000aab0f26&acdnat=1442234780_dfbadf711815831a2078ac4a70e25a66.
618. Vitriol VG, Ballesteros ST, Florenzano RU, Weil KP, Benadof DF. Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. *Psychiatric services* (Washington, DC). 2009;60(7):936-42.
619. Vrijzen JN, Fischer VS, Muller BW, Scherbaum N, Becker ES, Rinck M, et al. Cognitive bias modification as an add-on treatment in clinical depression: Results from a placebo-controlled, single-blinded randomized control trial. *Journal of affective disorders*. 2018;238:342-50.
620. Wagner B, Horn AB, Maercker A. Internet-based versus face-to-face cognitive-behavioral intervention for depression: a randomized controlled non-inferiority trial. *Journal of affective disorders*. 2014;152-154:113-21.
621. Walker JV, Lampropoulos GK. A comparison of self-help (homework) activities for mood enhancement: Results from a brief randomized controlled trial. *Journal of psychotherapy integration* [Internet]. 2014; 24(1):[46-64 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/401/CN-00986401/frame.html>.
622. Warmerdam L, Straten A, Twisk J, Riper H, Cuijpers P. Internet-based treatment for adults with depressive symptoms: randomized controlled trial. *Journal of medical Internet research* [Internet]. 2008; 10(4):[e44 p.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/612/CN-00667612/frame.html>.
623. Watkins ER, Baeyens CB, Read R. Concreteness training reduces dysphoria: proof-of-principle for repeated cognitive bias modification in depression. *Journal of abnormal psychology*. 2009;118(1):55-64.
624. Watkins ER, Taylor RS, Byng R, Baeyens C, Read R, Pearson K, et al. Guided self-help concreteness training as an intervention for major depression in primary care: a Phase II randomized controlled trial. *Psychological medicine*. 2012;42(7):1359-71.
625. Watson JC, Gordon LB, Stermac L, Kalogerakos F, Steckley P. Comparing the effectiveness of process-experiential with cognitive-behavioral psychotherapy in the treatment of depression. *Journal of consulting and clinical psychology*. 2003;71(4):773-81.
626. Watt LM, Cappeliez P. Integrative and instrumental reminiscence therapies for depression in older adults: Intervention strategies and treatment effectiveness. *Aging & mental health*. 2000;4(2):166-77.
627. Watts S, Mackenzie A, Thomas C, Griskaitis A, Mewton L, Williams A, et al. CBT for depression: a pilot RCT comparing mobile phone vs. computer. *BMC psychiatry*. 2013;13:49.
628. Weissman MM, Prusoff BA, Dimascio A, Neu C, Goklaney M, Klerman GL. The efficacy of drugs and psychotherapy in the treatment of acute depressive episodes. *The American journal of psychiatry*. 1979;136(4b):555-8.
629. Werner-Seidler A, Hitchcock C, Bevan A, McKinnon A, Gillard J, Dahm T, et al. A cluster randomized controlled platform trial comparing group MEMory specificity training (MEST) to group psychoeducation and supportive counselling (PSC) in the treatment of recurrent depression. *Behaviour research and therapy*. 2018;105:1-9.
630. Wickberg B, Hwang CP. Counselling of postnatal depression: a controlled study on a population based Swedish sample. *Journal of affective disorders*. 1996;39(3):209-16.
631. Wiersma JE, Schaik DJF, Hoogendorn AW, Dekker JJ, Van HL, Schoevers RA, et al. The effectiveness of the cognitive behavioral analysis system of psychotherapy for chronic depression: A randomized controlled trial. *Psychotherapy and psychosomatics* [Internet]. 2014; 83(5):[263-9 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/826/CN-01001826/frame.html>.
632. Wierzbicki M, Bartlett TS. The efficacy of group and individual cognitive therapy for mild depression. *Cognitive Therapy and Research*. 1987;11(3):337-42.
633. Wiklund I, Mohlkert P, Edman G. Evaluation of a brief cognitive intervention in patients with signs of postnatal depression: a randomized controlled trial. *Acta obstetricia et gynecologica Scandinavica*. 2010;89(8):1100-4.

634. Wiles N, Thomas L, Abel A, Ridgway N, Turner N, Campbell J, et al. Cognitive behavioural therapy as an adjunct to pharmacotherapy for primary care based patients with treatment resistant depression: results of the CoBaIT randomised controlled trial. *Lancet*. 2013;381(9864):375-84.
635. Wiles NJ, Hollinghurst S, Mason V, Musa M, Burt V, Hyde J, et al. A randomized controlled trial of cognitive behavioural therapy as an adjunct to pharmacotherapy in primary care based patients with treatment resistant depression: A pilot study. *Behavioural and Cognitive Psychotherapy*. 2008;36(1):21-33.
636. Williams AD, Blackwell SE, Mackenzie A, Holmes EA, Andrews G. Combining imagination and reason in the treatment of depression: A randomized controlled trial of internet-based cognitive-bias modification and internet-CBT for depression. *Journal of consulting and clinical psychology* [Internet]. 2013; 81(5):[793-9 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/389/CN-00974389/frame.html>
http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3780629/pdf/ccp_81_5_793.pdf.
637. Williams AD, O'Moore K, Blackwell SE, Smith J, Holmes EA, Andrews G. Positive imagery cognitive bias modification (CBM) and internet-based cognitive behavioral therapy (iCBT): a randomized controlled trial. *Journal of affective disorders*. 2015;178:131-41.
638. Williams C, McClay CA, Matthews L, McConnachie A, Haig C, Walker A, et al. Community-based group guided self-help intervention for low mood and stress: Randomised controlled trial. *British Journal of Psychiatry*. 2018;212(2):88-95.
639. Williams C, Wilson P, Morrison J, McMahan A, Andrew W, Allan L, et al. Guided Self-Help Cognitive Behavioural Therapy for Depression in Primary Care: A Randomised Controlled Trial. *PLoS ONE*. 2013;8(1).
640. Williams JW, Barrett J, Oxman T, Frank E, Katon W, Sullivan M, et al. Treatment of dysthymia and minor depression in primary care: A randomized controlled trial in older adults. *Jama* [Internet]. 2000; 284(12):[1519-26 pp.]. Available from:
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/681/CN-00299681/frame.html>
<http://jama.jamanetwork.com/data/Journals/JAMA/4753/JOC00569.pdf>.
641. Wilson GL. Psychotherapy with depressed incarcerated felons: a comparative evaluation of treatments. *Psychological reports*. 1990;67(3 Pt 1):1027-41.
642. Wilson PH. Combined pharmacological and behavioural treatment of depression. *Behaviour Research and Therapy*. 1982;20(2):173-84.
643. Wilson PH, Goldin JC, Charbonneau-Powis M. Comparative efficacy of behavioral and cognitive treatments of depression. *Cognitive therapy and research*. 1983;7(2):111-24.
644. Wollersheim JP, Wilson GL. Group treatment of unipolar depression: A comparison of coping, supportive, bibliotherapy, and delayed treatment groups. *Professional Psychology: Research and Practice*. 1991;22(6):496.
645. Wong DF. Cognitive and health-related outcomes of group cognitive behavioural treatment for people with depressive symptoms in Hong Kong: randomized wait-list control study. *The Australian and New Zealand journal of psychiatry*. 2008;42(8):702-11.
646. Wong DF. Cognitive behavioral treatment groups for people with chronic depression in Hong Kong: a randomized wait-list control design. *Depression and anxiety*. 2008;25(2):142-8.
647. Wong SYS, Sun YY, Chan ATY, Leung MKW, Chao DVK, Li CCK, et al. Treating Subthreshold Depression in Primary Care: a Randomized Controlled Trial of Behavioral Activation With Mindfulness. *Annals of family medicine*. 2018;16(2):111-9.
648. Wright JH, Wright AS, Albano AM, Basco MR, Goldsmith LJ, Raffield T, et al. Computer-assisted cognitive therapy for depression: maintaining efficacy while reducing therapist time. *The American journal of psychiatry*. 2005;162(6):1158-64.
649. Wuthrich VM, Rapee RM. Randomised controlled trial of group cognitive behavioural therapy for comorbid anxiety and depression in older adults. *Behaviour research and therapy* [Internet]. 2013; 51(12):[779-86 pp.]. Available from: <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/045/CN-00915045/frame.html>
http://ac.els-cdn.com/S0005796713001563/1-s2.0-S0005796713001563-main.pdf?_tid=58f675c4-5adf-11e5-8487-00000aacb35f&acdnat=1442235282_2dfb7de052fd54dfc45aa82ce68f0d01.
650. Wuthrich VM, Rapee RM, Kangas M, Perini S. Randomized controlled trial of group cognitive behavioral therapy compared to a discussion group for co-morbid anxiety and depression in older adults. *Psychological medicine*. 2016;46(4):785-95.
651. Yang TT, Hsiao FH, Wang KC, Ng SM, Ho RT, Chan CL, et al. The effect of psychotherapy added to pharmacotherapy on cortisol responses in outpatients with major depressive disorder. *The Journal of nervous and mental disease*. 2009;197(6):401-6.

652. Yang W, Ding Z, Dai T, Peng F, Zhang JX. Attention Bias Modification training in individuals with depressive symptoms: A randomized controlled trial. *Journal of behavior therapy and experimental psychiatry* [Internet]. 2015; 49:[101-11 pp.]. Available from:
<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-3113.2014.00475.x>
http://ac.els-cdn.com/S0005791614000755/1-s2.0-S0005791614000755-main.pdf?_tid=74fcd4d6-b60e-11e5-bad8-00000aab0f6b&acdnat=1452261071_44276b0335c97a99c9659ecff10ef4c3.
653. Yang X, Zhao J, Chen Y, Zu S, Zhao J. Comprehensive self-control training benefits depressed college students: a six-month randomized controlled intervention trial. *Journal of affective disorders*. 2018;226:251-60.
654. Yeung A, Wang F, Feng F, Zhang J, Cooper A, Hong L, et al. Outcomes of an online computerized cognitive behavioral treatment program for treating chinese patients with depression: A pilot study. *Asian Journal of Psychiatry*. 2017.
655. Yiend J, Lee JS, Tekes S, Atkins L, Mathews A, Vrinten M, et al. Modifying interpretation in a clinically depressed sample using 'cognitive bias modification-errors': A double blind randomised controlled trial. *Cognitive Therapy and Research*. 2014;38(2):146-59.
656. Zagorscak P, Heinrich M, Sommer D, Wagner B, Knaevelsrud C. Benefits of Individualized Feedback in Internet-Based Interventions for Depression: a Randomized Controlled Trial. *Psychotherapy and psychosomatics*. 2018;87(1):32-45.
657. Zemestani M, Davoodi I, Honarmand MM, Zargar Y, Ottaviani C. Comparative effects of group metacognitive therapy versus behavioural activation in moderately depressed students. *Journal of Mental Health*. 2016;25(6):479-85.
658. Zettle RD, Rains JC. Group cognitive and contextual therapies in treatment of depression. *Journal of clinical psychology*. 1989.
659. Zhu S, Zhu K, Jiang D, Shi J. Social cognition and interaction training for major depression: A preliminary study. *Psychiatry Research*. 2018;270:890-4.
660. Zisook S, Peterkin J, Goggin KJ, Sledge P, Atkinson JH, Grant I. Treatment of major depression in HIV-seropositive men. HIV Neurobehavioral Research Center Group. *The Journal of clinical psychiatry*. 1998;59(5):217-24.
661. Zu S, Xiang Y-T, Liu J, Zhang L, Wang G, Ma X, et al. A comparison of cognitive-behavioral therapy, antidepressants, their combination and standard treatment for Chinese patients with moderate-severe major depressive disorders. *Journal of Affective Disorders*. 2014;152-154:262-7.